

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

エネルギー(kcal)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	1,930	520	58	925	1,011	1,349	1,526	1,825	2,288	2,740	2,898	3,135
	30-39歳	58	2,244	726	90	647	1,139	1,294	1,746	2,172	2,704	3,231	3,701	4,135
	40-49歳	96	2,076	524	56	877	1,196	1,401	1,699	2,061	2,396	2,790	2,964	3,473
	50-59歳	84	2,144	511	60	905	1,432	1,530	1,774	2,113	2,447	2,798	3,108	3,871
	60-69歳	115	2,227	467	45	1,069	1,456	1,580	1,934	2,193	2,518	2,890	3,109	3,224
	70歳以上	168	2,078	524	52	991	1,269	1,432	1,678	2,109	2,390	2,756	3,088	3,370
	(再掲)20歳以上※	563	2,112	550	28	950	1,245	1,457	1,708	2,118	2,437	2,818	3,113	3,596
	(再掲)65-74歳	130	2,211	467	46	995	1,455	1,584	1,908	2,185	2,468	2,884	3,014	3,384
	(再掲)75歳以上	107	2,023	504	62	1,023	1,245	1,372	1,649	2,009	2,348	2,662	2,961	3,341
	(再掲)20-49歳※	196	2,074	597	37	883	1,193	1,349	1,678	2,071	2,431	2,820	3,108	3,944
(再掲)40-74歳※	356	2,154	509	30	1,000	1,386	1,532	1,807	2,137	2,455	2,827	3,075	3,392	
女性	20-29歳	40	1,548	412	84	610	837	907	1,326	1,540	1,840	1,994	2,304	2,395
	30-39歳	87	1,657	396	53	637	1,028	1,237	1,410	1,629	1,875	2,177	2,273	3,304
	40-49歳	85	1,781	406	48	933	1,072	1,251	1,462	1,804	2,119	2,310	2,332	3,115
	50-59歳	97	1,732	348	30	956	1,155	1,273	1,547	1,728	1,931	2,079	2,389	2,912
	60-69歳	145	1,759	476	41	613	1,053	1,240	1,500	1,679	1,987	2,342	2,733	3,155
	70歳以上	187	1,642	390	30	772	973	1,137	1,376	1,670	1,910	2,134	2,263	2,577
	(再掲)20歳以上※	641	1,684	412	21	757	989	1,169	1,429	1,676	1,928	2,214	2,344	2,915
	(再掲)65-74歳	151	1,746	447	34	749	1,120	1,307	1,431	1,680	1,968	2,304	2,578	3,144
	(再掲)75歳以上	137	1,614	405	36	743	921	1,049	1,347	1,666	1,898	2,125	2,310	2,738
	(再掲)20-49歳※	212	1,668	415	38	643	928	1,098	1,397	1,655	1,936	2,229	2,315	2,823
(再掲)40-74歳※	377	1,752	406	21	841	1,116	1,292	1,471	1,721	1,977	2,271	2,412	3,067	
男女計	20-29歳	82	1,746	506	58	629	927	1,019	1,438	1,706	2,050	2,425	2,773	3,135
	30-39歳	145	1,956	655	61	641	1,098	1,241	1,504	1,802	2,229	2,962	3,235	4,097
	40-49歳	181	1,931	491	46	935	1,153	1,307	1,612	1,926	2,237	2,660	2,803	3,128
	50-59歳	181	1,938	483	43	967	1,258	1,426	1,641	1,865	2,221	2,518	2,904	3,578
	60-69歳	260	1,990	526	40	774	1,173	1,396	1,597	1,948	2,339	2,712	2,965	3,214
	70歳以上	355	1,823	499	36	836	1,042	1,217	1,496	1,760	2,132	2,450	2,740	3,322
	(再掲)20歳以上※	1,204	1,891	529	23	839	1,083	1,263	1,532	1,839	2,195	2,593	2,863	3,304
	(再掲)65-74歳	281	1,963	512	33	798	1,254	1,367	1,590	1,912	2,304	2,643	2,910	3,257
	(再掲)75歳以上	244	1,773	488	42	790	1,010	1,155	1,461	1,720	2,111	2,393	2,645	3,298
	(再掲)20-49歳※	408	1,876	555	35	778	1,027	1,236	1,506	1,812	2,178	2,645	2,896	3,548
(再掲)40-74歳※	733	1,952	502	24	912	1,197	1,374	1,601	1,910	2,247	2,636	2,900	3,222	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

総たんぱく質(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	71.4	24.4	3.90	31.6	35.4	40.0	55.2	66.3	92.2	106.7	110.5	129.8
	30-39歳	58	82.9	28.2	3.83	23.9	36.2	51.4	65.0	78.4	102.2	126.4	139.0	143.1
	40-49歳	96	73.0	20.3	2.30	25.1	42.5	44.7	58.2	74.2	88.9	99.1	104.4	140.2
	50-59歳	84	75.9	22.1	2.48	27.2	45.7	50.1	62.2	73.4	85.0	104.6	116.2	167.0
	60-69歳	115	80.0	21.3	1.87	32.0	47.4	54.8	64.5	78.4	92.5	108.2	120.6	139.6
	70歳以上	168	76.5	21.6	2.01	27.0	39.9	48.1	59.3	78.8	94.0	101.4	111.3	124.8
	(再掲)20歳以上※	563	76.5	23.0	1.34	29.7	40.9	47.5	60.3	74.6	91.9	104.8	116.8	140.0
	(再掲)65-74歳	130	81.5	20.3	1.91	35.7	47.5	55.8	67.3	80.0	92.7	107.7	118.2	135.5
	(再掲)75歳以上	107	75.0	21.9	2.79	24.2	38.9	48.0	57.6	76.7	94.1	100.5	106.0	132.4
	(再掲)20-49歳※	196	75.3	24.5	2.20	28.0	38.7	44.3	58.4	73.0	91.7	105.5	122.5	141.6
(再掲)40-74歳※	356	76.7	21.3	1.10	32.0	44.4	49.1	62.0	76.1	91.0	103.1	113.5	139.5	
女性	20-29歳	40	60.8	20.7	3.98	23.4	26.5	36.2	47.2	57.9	74.7	93.1	101.2	110.1
	30-39歳	87	60.9	16.5	1.91	22.5	34.5	39.9	49.3	59.3	71.9	80.5	93.8	101.1
	40-49歳	85	61.6	15.9	1.54	30.6	37.3	40.4	50.5	60.6	69.1	83.7	92.8	110.2
	50-59歳	97	65.0	18.1	2.15	27.7	39.9	42.8	50.0	66.0	74.0	88.8	96.2	123.2
	60-69歳	145	69.1	21.6	1.80	18.2	36.8	41.7	53.9	67.6	83.9	96.2	105.2	130.9
	70歳以上	187	63.7	18.7	1.54	24.8	36.0	40.0	50.6	62.0	75.0	88.8	99.1	120.4
	(再掲)20歳以上※	641	63.8	18.9	0.92	25.2	35.7	40.5	50.6	61.5	74.6	88.9	99.1	114.9
	(再掲)65-74歳	151	68.9	21.2	1.86	18.4	36.3	46.0	53.2	69.0	82.3	95.4	104.0	130.8
	(再掲)75歳以上	137	62.8	18.4	1.39	25.5	36.0	38.3	49.8	61.1	73.4	86.2	101.1	113.4
	(再掲)20-49歳※	212	61.1	17.7	1.44	23.8	34.7	40.3	50.0	59.5	71.6	85.3	98.5	106.2
(再掲)40-74歳※	377	65.5	19.1	0.95	26.4	37.3	42.9	51.3	64.8	76.7	90.1	99.1	124.9	
男女計	20-29歳	82	66.3	23.2	3.44	23.9	34.0	39.5	49.4	61.3	78.0	102.5	110.3	129.8
	30-39歳	145	72.1	25.6	2.59	23.8	35.3	42.8	53.4	69.7	87.5	104.6	126.3	142.3
	40-49歳	181	67.4	19.1	1.50	31.5	39.6	44.3	51.6	65.2	80.0	94.6	101.6	116.6
	50-59歳	181	70.5	20.8	1.97	27.6	41.7	46.2	56.6	69.4	81.4	95.3	108.7	141.9
	60-69歳	260	74.5	22.1	1.60	25.1	40.8	47.6	58.8	72.7	88.8	102.1	116.0	132.8
	70歳以上	355	69.0	20.9	1.51	25.3	36.8	43.2	53.7	66.9	82.3	97.7	103.3	124.3
	(再掲)20歳以上※	1,204	69.9	21.9	1.01	26.4	37.5	43.4	54.3	67.7	82.9	99.5	108.3	132.3
	(再掲)65-74歳	281	74.8	21.7	1.63	21.2	41.8	48.9	59.3	73.2	89.0	100.8	115.5	131.9
	(再掲)75歳以上	244	67.5	20.7	1.70	25.4	36.3	40.1	52.9	65.2	81.7	96.5	102.7	121.8
	(再掲)20-49歳※	408	68.4	22.6	1.69	24.9	35.5	41.6	52.4	65.4	80.1	101.3	110.2	137.8
(再掲)40-74歳※	733	71.1	20.9	0.83	27.7	40.5	46.3	56.4	69.4	84.2	98.4	106.7	131.6	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

たんぱく質エネルギー比率(%)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	14.8	3.1	0.70	8.8	10.9	11.3	12.5	14.6	15.8	20.2	20.7	22.4
	30-39歳	58	15.0	3.4	0.67	9.5	10.3	10.8	12.7	14.5	16.6	17.8	22.1	25.0
	40-49歳	96	14.2	2.6	0.32	8.7	10.4	10.8	12.1	13.9	15.8	17.9	18.8	21.2
	50-59歳	84	14.2	2.7	0.29	7.9	10.4	11.4	12.1	14.1	15.7	17.6	19.6	22.7
	60-69歳	115	14.4	2.7	0.27	9.3	10.5	10.7	12.1	14.4	16.5	18.4	18.8	19.9
	70歳以上	168	14.8	2.7	0.20	7.2	10.4	11.6	13.1	14.9	16.8	18.3	19.6	21.3
	(再掲)20歳以上※	563	14.6	2.8	0.24	8.8	10.5	11.1	12.4	14.4	16.4	18.3	19.7	22.6
	(再掲)65-74歳	130	14.8	2.6	0.26	9.6	10.6	11.4	12.8	14.7	16.8	18.5	19.2	20.7
	(再掲)75歳以上	107	14.9	2.8	0.25	6.4	10.3	11.2	13.1	15.0	16.9	18.3	19.6	21.3
	(再掲)20-49歳※	196	14.6	3.0	0.41	8.8	10.6	11.1	12.4	14.3	16.2	18.2	20.7	24.1
(再掲)40-74歳※	356	14.3	2.6	0.15	8.8	10.6	11.1	12.1	14.3	16.2	18.2	18.8	20.9	
女性	20-29歳	40	15.8	3.6	0.80	10.2	10.8	11.2	13.0	15.2	17.7	21.5	22.3	22.8
	30-39歳	87	14.8	2.5	0.24	9.5	11.5	11.9	13.0	14.3	16.1	18.7	20.3	21.3
	40-49歳	85	13.9	2.5	0.25	10.3	10.7	11.2	12.0	13.8	15.6	17.4	18.2	23.1
	50-59歳	97	15.0	3.1	0.48	8.8	10.8	11.1	12.6	14.9	17.0	18.5	20.9	24.5
	60-69歳	145	15.7	2.8	0.26	9.5	11.6	12.1	13.4	15.6	17.5	19.7	20.9	23.1
	70歳以上	187	15.5	2.8	0.24	8.2	11.7	12.3	13.6	15.0	17.1	19.7	20.7	22.4
	(再掲)20歳以上※	641	15.2	3.0	0.20	9.8	11.1	11.6	13.0	14.7	17.0	19.5	20.7	22.9
	(再掲)65-74歳	151	15.8	3.0	0.30	8.0	11.3	12.3	13.6	15.6	17.5	20.2	20.8	23.0
	(再掲)75歳以上	137	15.6	2.6	0.23	10.3	12.0	12.6	13.7	14.9	17.1	19.7	20.3	23.2
	(再掲)20-49歳※	212	14.8	3.0	0.33	10.1	10.8	11.3	12.4	14.3	16.7	19.4	21.0	22.7
(再掲)40-74歳※	377	15.0	2.9	0.22	9.1	10.9	11.4	12.6	14.7	16.9	18.9	20.7	23.1	
男女計	20-29歳	82	15.3	3.4	0.65	8.9	10.9	11.2	12.8	14.9	17.4	20.7	21.8	22.8
	30-39歳	145	14.9	3.0	0.39	9.5	10.7	11.5	12.8	14.4	16.5	17.9	20.4	24.5
	40-49歳	181	14.1	2.5	0.21	8.8	10.7	11.0	12.1	13.8	15.6	17.5	18.5	21.5
	50-59歳	181	14.6	2.9	0.35	8.5	10.8	11.3	12.1	14.5	16.5	18.0	19.7	23.6
	60-69歳	260	15.1	2.8	0.22	9.5	10.7	11.4	12.7	14.9	17.1	18.7	20.0	22.5
	70歳以上	355	15.2	2.8	0.21	8.1	11.1	11.9	13.4	14.9	16.9	19.3	20.2	22.2
	(再掲)20歳以上※	1,204	14.9	2.9	0.20	9.2	10.7	11.4	12.7	14.6	16.8	18.8	20.4	22.8
	(再掲)65-74歳	281	15.3	2.9	0.25	9.5	10.8	11.7	13.3	15.1	17.2	19.3	20.3	22.5
	(再掲)75歳以上	244	15.3	2.7	0.19	8.5	11.2	12.1	13.6	15.0	17.0	19.2	20.2	21.9
	(再掲)20-49歳※	408	14.7	3.0	0.33	9.4	10.7	11.2	12.4	14.3	16.4	18.9	20.7	23.1
(再掲)40-74歳※	733	14.7	2.8	0.17	8.9	10.7	11.3	12.4	14.5	16.6	18.3	19.7	22.7	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

動物性たんぱく質(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	42.1	22.3	3.52	13.3	14.0	16.2	27.0	36.0	55.2	81.3	86.3	95.4
	30-39歳	58	47.2	22.5	3.53	11.8	14.4	19.3	30.1	44.0	60.9	80.8	90.5	110.7
	40-49歳	96	39.5	17.2	1.68	3.2	12.4	16.2	25.3	39.0	51.1	60.7	75.1	95.4
	50-59歳	84	39.4	16.8	1.94	11.3	16.7	18.6	26.1	39.7	45.1	60.8	75.1	104.7
	60-69歳	115	41.9	18.6	1.64	6.9	15.1	18.4	29.3	40.9	53.3	69.6	78.8	88.0
	70歳以上	168	41.0	17.9	1.53	4.4	14.1	20.5	26.6	39.7	53.9	65.1	77.3	88.6
	(再掲)20歳以上※	563	41.6	19.2	1.22	9.7	14.4	18.6	27.4	39.8	53.1	68.6	80.6	94.2
	(再掲)65-74歳	130	43.7	17.8	1.67	6.5	15.4	21.9	31.8	42.9	54.8	69.5	78.6	85.9
	(再掲)75歳以上	107	40.6	18.9	2.12	0.5	12.3	20.5	25.4	38.1	54.8	66.2	77.7	92.7
	(再掲)20-49歳※	196	42.6	20.7	2.04	10.1	13.9	17.0	27.4	39.9	54.6	76.4	85.9	95.6
(再掲)40-74歳※	356	40.5	17.4	0.93	9.0	14.2	18.6	27.4	40.4	50.4	64.3	75.6	88.0	
女性	20-29歳	40	35.1	19.0	3.60	3.8	7.1	12.1	23.8	32.1	43.0	63.3	75.7	81.7
	30-39歳	87	32.9	13.3	1.48	6.3	14.6	15.5	23.2	31.5	41.3	51.7	58.3	66.5
	40-49歳	85	32.0	13.6	1.08	2.1	15.0	18.5	21.7	29.3	38.4	50.2	56.5	74.6
	50-59歳	97	34.8	15.5	1.95	6.7	11.1	14.7	24.4	33.4	43.7	52.6	68.3	83.7
	60-69歳	145	37.0	16.7	1.40	6.1	12.0	16.4	24.5	36.7	48.7	58.5	68.3	80.5
	70歳以上	187	34.0	15.5	1.36	4.3	13.6	16.2	21.8	33.1	42.2	55.2	67.7	77.8
	(再掲)20歳以上※	641	34.3	15.7	0.88	5.4	12.2	16.1	22.9	32.5	42.9	54.3	66.5	81.5
	(再掲)65-74歳	151	37.9	17.0	1.45	1.8	13.2	18.5	25.6	36.4	48.7	62.1	72.0	80.5
	(再掲)75歳以上	137	33.1	14.6	1.16	6.5	13.2	15.4	20.9	32.8	42.3	51.8	64.0	77.5
	(再掲)20-49歳※	212	33.3	15.5	1.44	4.2	12.1	15.7	23.0	30.7	40.2	52.9	65.0	79.9
(再掲)40-74歳※	377	34.9	15.8	0.88	6.6	13.6	17.2	23.3	32.5	43.5	54.5	68.2	78.1	
男女計	20-29歳	82	38.7	20.9	3.13	4.1	12.1	14.9	25.1	34.4	49.7	77.8	83.5	95.4
	30-39歳	145	40.2	19.9	2.38	9.6	14.7	18.0	24.2	37.7	50.3	69.1	80.7	101.0
	40-49歳	181	35.8	16.0	1.12	3.5	13.4	18.2	24.1	34.0	45.2	55.9	66.1	83.1
	50-59歳	181	37.1	16.3	1.66	7.5	13.7	18.2	25.8	36.2	44.9	57.2	69.2	95.4
	60-69歳	260	39.4	17.8	1.34	6.8	13.0	16.8	25.7	37.9	50.1	64.9	74.8	83.4
	70歳以上	355	36.9	16.9	1.22	5.4	13.8	17.0	23.5	35.3	45.9	59.1	70.1	80.4
	(再掲)20歳以上※	1,204	37.9	17.9	0.99	6.8	13.6	17.2	25.1	35.4	47.4	61.3	75.7	87.1
	(再掲)65-74歳	281	40.6	17.6	1.32	5.5	15.3	19.9	27.4	38.8	52.0	65.8	77.2	82.5
	(再掲)75歳以上	244	36.0	16.8	1.27	5.5	12.9	16.6	22.4	34.6	45.3	58.9	68.2	80.5
	(再掲)20-49歳※	408	38.0	18.9	1.67	5.7	13.2	16.8	24.5	34.6	47.5	65.5	80.1	91.0
(再掲)40-74歳※	733	37.7	16.8	0.79	7.5	13.7	18.0	25.6	36.1	47.2	58.8	70.9	82.8	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

動物性たんぱく質比率(%)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	56.4	13.3	1.95	25.9	35.4	40.2	44.7	55.8	66.0	76.6	77.8	78.9
	30-39歳	58	55.3	13.2	2.01	25.8	28.8	37.5	45.3	56.7	65.4	72.6	73.4	77.4
	40-49歳	96	52.1	12.8	1.35	8.3	28.9	33.7	44.3	55.8	61.2	66.3	73.0	79.2
	50-59歳	84	50.8	10.5	1.66	16.8	31.7	35.7	44.6	50.5	59.3	63.2	65.1	77.5
	60-69歳	115	50.6	13.1	1.16	15.5	25.7	31.4	44.0	50.8	61.6	67.2	70.4	76.6
	70歳以上	168	51.6	12.7	1.01	12.5	30.5	35.2	44.0	52.1	60.3	66.9	71.5	77.0
	(再掲)20歳以上※	563	52.6	12.8	0.80	18.9	29.8	36.0	44.6	53.3	61.6	68.0	73.5	77.9
	(再掲)65-74歳	130	52.0	12.2	1.14	15.6	26.5	36.4	45.3	52.6	61.2	66.9	69.6	74.3
	(再掲)75歳以上	107	51.8	13.8	1.39	2.0	30.1	35.1	43.5	52.1	61.2	69.6	75.2	78.0
	(再掲)20-49歳※	196	54.5	13.2	1.20	25.7	29.7	38.3	44.6	55.8	63.7	72.6	76.2	78.4
(再掲)40-74歳※	356	51.2	12.1	0.87	16.6	29.1	35.2	44.6	51.8	60.3	64.8	69.0	75.7	
女性	20-29歳	40	54.6	15.6	3.10	14.5	17.8	35.6	46.7	55.1	67.0	74.1	77.9	79.5
	30-39歳	87	52.8	11.8	1.45	18.9	33.7	39.2	42.9	52.1	64.1	68.0	69.1	75.4
	40-49歳	85	51.0	12.6	1.73	5.9	29.7	34.1	42.7	53.0	59.8	67.5	71.1	74.5
	50-59歳	97	51.9	13.6	1.63	19.2	26.0	31.0	45.3	54.2	59.9	70.3	75.1	77.4
	60-69歳	145	51.8	12.1	0.98	19.8	29.3	34.0	45.5	52.6	60.8	66.8	70.8	76.5
	70歳以上	187	51.5	13.1	1.09	10.4	31.6	34.7	42.4	53.3	60.2	68.5	70.9	78.7
	(再掲)20歳以上※	641	52.1	13.1	0.91	17.5	30.3	34.7	44.1	53.2	61.0	68.5	72.6	79.5
	(再掲)65-74歳	151	53.1	12.7	0.95	7.3	32.0	36.4	45.6	53.7	61.2	68.0	73.2	78.7
	(再掲)75歳以上	137	51.2	12.5	1.26	19.6	31.4	34.6	41.6	52.9	60.2	68.2	70.5	79.2
	(再掲)20-49歳※	212	52.7	13.5	1.60	15.4	30.4	35.9	44.0	53.3	63.9	69.0	72.8	78.5
(再掲)40-74歳※	377	51.7	13.0	0.84	19.8	28.1	33.8	44.3	53.2	60.1	67.5	72.4	77.9	
男女計	20-29歳	82	55.5	14.4	1.98	15.2	29.9	38.4	46.0	55.2	66.1	75.8	78.4	79.5
	30-39歳	145	54.1	12.5	1.59	22.6	32.7	38.3	44.0	54.4	64.3	70.0	73.3	76.8
	40-49歳	181	51.6	12.7	1.39	19.3	29.2	34.0	43.7	53.8	60.8	66.7	72.6	75.8
	50-59歳	181	51.4	12.1	1.47	19.2	27.9	34.3	45.3	52.8	59.6	65.0	71.1	77.5
	60-69歳	260	51.2	12.6	0.89	17.5	26.5	33.2	44.7	52.0	61.1	66.9	70.4	75.6
	70歳以上	355	51.6	12.9	0.89	12.9	31.5	34.9	43.2	52.8	60.2	67.8	70.8	78.5
	(再掲)20歳以上※	1,204	52.4	13.0	0.81	17.9	30.1	35.3	44.3	53.3	61.3	68.5	72.7	78.5
	(再掲)65-74歳	281	52.6	12.5	0.90	16.0	30.4	36.4	45.4	53.1	61.2	67.4	70.8	77.7
	(再掲)75歳以上	244	51.4	13.0	1.05	14.2	31.0	34.9	42.5	52.3	60.3	68.5	71.0	78.2
	(再掲)20-49歳※	408	53.6	13.3	1.28	17.8	30.4	37.1	44.4	54.4	63.7	70.2	74.0	79.1
(再掲)40-74歳※	733	51.5	12.5	0.81	19.0	29.0	34.6	44.5	52.6	60.2	66.3	70.8	77.4	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

脂質(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	58.6	21.1	2.28	18.3	21.6	31.8	43.0	58.1	74.2	91.0	99.2	103.4
	30-39歳	58	64.0	27.9	3.49	14.5	27.6	29.2	43.6	62.0	79.2	103.6	115.2	173.9
	40-49歳	96	60.7	22.6	2.15	17.2	26.0	33.2	43.3	59.6	76.8	88.3	99.0	134.1
	50-59歳	84	62.6	22.8	2.22	20.4	28.7	33.7	44.1	61.9	75.1	97.3	102.7	127.0
	60-69歳	115	60.6	23.0	2.79	21.9	31.4	35.8	43.2	55.7	74.4	92.7	114.6	125.9
	70歳以上	168	52.5	23.0	2.26	8.1	16.8	24.1	35.8	51.1	65.6	82.3	91.3	123.7
	(再掲)20歳以上※	563	59.3	23.5	1.25	15.7	25.0	31.3	42.4	57.4	73.7	89.7	102.1	127.2
	(再掲)65-74歳	130	58.3	21.4	2.24	15.5	25.5	31.5	42.4	55.8	73.4	85.1	94.4	114.7
	(再掲)75歳以上	107	50.7	22.2	2.85	7.4	15.9	22.7	33.4	50.1	62.2	79.3	91.1	131.8
	(再掲)20-49歳※	196	60.9	23.8	1.59	18.0	25.3	31.3	43.6	59.6	76.3	90.2	102.6	131.1
(再掲)40-74歳※	356	60.5	23.0	1.40	16.9	28.0	33.7	43.3	59.0	74.9	91.2	103.0	127.3	
女性	20-29歳	40	48.9	15.3	2.26	17.7	20.7	32.4	38.6	47.0	57.8	67.1	81.3	95.0
	30-39歳	87	51.3	17.3	2.65	10.2	24.9	29.4	38.0	49.3	64.0	73.2	79.1	99.6
	40-49歳	85	55.4	20.9	2.12	13.6	26.2	30.9	38.5	54.6	72.0	81.4	90.4	134.8
	50-59歳	97	55.1	20.1	1.74	17.6	27.4	33.3	40.8	53.4	66.6	81.4	101.3	113.7
	60-69歳	145	51.8	24.7	2.25	6.9	21.7	27.9	34.7	48.4	62.2	75.3	100.1	144.8
	70歳以上	187	45.3	17.9	1.48	8.6	19.2	23.2	32.1	44.5	57.7	67.7	75.7	97.5
	(再掲)20歳以上※	641	50.5	19.9	0.93	11.4	22.0	27.7	37.0	48.1	61.7	73.7	83.9	112.5
	(再掲)65-74歳	151	50.9	23.9	1.97	7.9	21.9	27.8	35.2	47.5	60.5	76.0	100.0	143.5
	(再掲)75歳以上	137	44.1	18.1	1.64	7.4	16.9	22.7	30.4	42.8	56.2	67.2	75.7	95.0
	(再掲)20-49歳※	212	52.0	18.3	1.33	14.8	24.1	30.9	38.5	49.3	64.8	77.2	84.4	95.4
(再掲)40-74歳※	377	53.2	21.6	1.12	13.7	25.1	30.0	37.9	49.4	65.3	80.2	91.9	127.3	
男女計	20-29歳	82	53.9	19.1	1.84	17.7	21.0	32.5	41.3	51.8	65.8	79.1	95.9	103.4
	30-39歳	145	57.7	24.1	2.38	12.9	27.2	29.4	41.3	54.3	71.7	84.4	104.5	145.4
	40-49歳	181	58.1	21.9	1.77	15.8	26.2	31.3	42.3	55.9	73.1	85.8	92.3	131.5
	50-59歳	181	58.8	21.8	1.50	20.1	28.3	33.7	43.1	55.4	70.4	89.1	102.7	119.9
	60-69歳	260	56.1	24.2	1.95	11.9	26.2	30.5	39.3	51.9	69.8	85.9	104.3	134.4
	70歳以上	355	48.3	20.5	1.51	8.7	18.6	23.2	33.4	46.7	60.5	75.0	84.3	113.3
	(再掲)20歳以上※	1,204	54.7	22.2	0.99	13.7	23.0	29.3	38.5	52.2	67.1	82.0	95.0	119.4
	(再掲)65-74歳	281	54.4	23.0	1.80	12.6	23.5	29.3	37.9	51.5	65.4	83.2	96.6	130.2
	(再掲)75歳以上	244	46.7	20.0	1.84	8.1	16.8	22.8	32.3	46.2	59.9	70.5	81.0	96.9
	(再掲)20-49歳※	408	56.6	21.7	1.27	17.0	25.3	31.0	41.7	54.7	70.3	82.3	95.0	117.0
(再掲)40-74歳※	733	56.8	22.6	1.09	15.8	26.4	31.2	40.4	53.9	70.4	85.8	99.7	126.9	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

脂肪エネルギー比率(%)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	27.1	6.0	0.47	12.2	14.8	18.4	24.0	27.3	29.8	34.9	38.5	40.0
	30-39歳	58	25.3	6.0	0.72	11.8	17.7	18.4	21.4	23.9	28.4	32.9	35.2	48.0
	40-49歳	96	26.3	7.0	0.78	11.5	14.4	16.5	21.0	25.9	31.7	37.1	38.7	41.9
	50-59歳	84	26.0	6.2	0.68	12.8	13.9	17.6	21.3	26.6	30.0	32.8	35.1	43.9
	60-69歳	115	24.3	6.8	0.89	10.5	14.5	15.5	20.0	24.3	28.2	32.1	32.8	50.5
	70歳以上	168	22.4	7.5	0.64	6.1	9.6	13.0	17.4	22.2	25.9	33.0	36.6	46.7
	(再掲)20歳以上※	563	25.0	6.9	0.35	9.0	14.0	16.1	20.6	25.1	29.3	33.0	37.1	43.5
	(再掲)65-74歳	130	23.6	6.9	0.72	8.6	12.8	14.8	18.5	23.6	28.2	32.2	35.0	43.0
	(再掲)75歳以上	107	22.2	7.4	0.85	4.9	8.9	12.5	17.5	22.2	26.0	32.0	36.2	45.9
	(再掲)20-49歳※	196	26.3	6.4	0.47	11.8	15.2	18.0	21.7	26.2	29.9	34.8	38.0	41.6
(再掲)40-74歳※	356	25.1	6.9	0.43	11.2	14.0	15.9	20.3	25.2	29.7	33.1	37.1	43.9	
女性	20-29歳	40	28.9	6.7	1.11	11.1	18.1	21.2	25.1	27.6	34.0	37.2	41.1	46.7
	30-39歳	87	27.7	6.6	0.93	9.3	18.3	19.6	22.0	27.7	32.1	35.9	38.6	48.6
	40-49歳	85	27.7	7.1	0.66	9.3	17.1	19.0	22.3	28.5	33.4	36.7	37.6	43.3
	50-59歳	97	28.4	7.8	0.68	12.6	17.5	19.4	22.6	27.1	32.4	37.6	42.8	53.4
	60-69歳	145	25.8	7.6	0.72	8.9	14.9	16.1	20.4	25.3	30.0	36.7	41.3	44.3
	70歳以上	187	24.5	7.0	0.59	8.5	13.2	15.4	20.0	24.3	28.7	33.7	36.6	41.6
	(再掲)20歳以上※	641	26.7	7.3	0.38	9.6	15.4	17.5	21.7	26.5	31.7	36.4	38.4	45.4
	(再掲)65-74歳	151	25.6	7.3	0.70	9.4	14.7	15.7	20.7	25.3	29.4	34.1	41.1	44.2
	(再掲)75歳以上	137	24.2	7.1	0.74	7.7	12.7	15.3	19.5	23.7	29.0	34.8	36.6	38.9
	(再掲)20-49歳※	212	28.1	6.8	0.55	9.8	17.8	19.1	23.5	28.1	33.3	36.4	37.9	46.3
(再掲)40-74歳※	377	26.9	7.4	0.37	10.5	15.6	17.5	21.9	26.8	31.7	36.9	39.8	45.0	
男女計	20-29歳	82	27.9	6.4	0.62	11.1	16.0	20.2	24.3	27.7	31.9	36.9	38.2	46.7
	30-39歳	145	26.5	6.4	0.62	10.9	18.0	19.1	21.6	25.9	30.9	34.7	36.6	48.3
	40-49歳	181	27.0	7.1	0.53	10.3	14.6	17.6	21.0	26.9	32.1	36.7	38.2	41.3
	50-59歳	181	27.2	7.1	0.55	12.8	16.4	18.8	22.1	27.0	30.9	36.5	39.7	53.0
	60-69歳	260	25.1	7.3	0.60	10.4	14.8	16.0	20.1	24.7	29.2	33.3	40.9	44.8
	70歳以上	355	23.6	7.2	0.54	7.7	12.3	14.5	18.9	23.3	28.0	33.4	36.6	42.1
	(再掲)20歳以上※	1,204	25.9	7.1	0.34	9.6	14.5	17.0	21.1	25.8	30.2	35.4	37.7	44.7
	(再掲)65-74歳	281	24.6	7.2	0.64	9.2	14.1	15.3	19.7	24.5	28.7	32.9	38.6	43.5
	(再掲)75歳以上	244	23.4	7.3	0.70	7.2	10.9	14.4	18.9	22.8	27.8	33.5	36.5	38.9
	(再掲)20-49歳※	408	27.1	6.7	0.42	11.2	16.5	19.0	22.7	27.2	31.8	36.0	38.0	44.6
(再掲)40-74歳※	733	26.0	7.2	0.33	11.2	14.6	16.7	21.0	25.8	30.5	35.7	38.4	44.0	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

動物性脂質(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	27.3	13.3	2.07	4.6	7.0	8.1	16.3	27.5	35.6	48.8	50.6	51.9
	30-39歳	58	35.7	19.5	2.33	7.7	10.7	16.9	22.3	34.2	42.5	61.3	68.9	134.7
	40-49歳	96	31.2	17.9	1.40	0.1	7.7	8.9	17.7	28.6	41.6	55.0	67.0	91.7
	50-59歳	84	31.5	15.0	1.66	9.1	12.6	14.9	18.8	29.5	44.0	54.9	59.5	67.4
	60-69歳	115	30.3	15.3	1.78	3.2	8.1	11.1	19.1	29.3	38.5	56.0	60.3	73.6
	70歳以上	168	27.9	14.8	1.43	1.7	5.5	9.2	17.6	25.8	38.5	49.3	55.7	60.7
	(再掲)20歳以上※	563	30.3	16.1	0.78	2.8	8.3	10.9	18.5	28.7	40.3	50.2	59.7	73.2
	(再掲)65-74歳	130	30.4	15.1	1.54	2.9	8.1	10.9	19.1	29.3	38.6	52.8	60.0	71.7
	(再掲)75歳以上	107	27.5	14.9	1.85	1.3	4.9	8.8	17.5	24.4	38.4	48.5	55.9	62.6
	(再掲)20-49歳※	196	31.2	17.2	1.13	2.0	8.2	10.2	18.7	28.9	41.1	49.8	62.4	91.4
(再掲)40-74歳※	356	30.6	16.0	0.83	2.8	8.3	12.1	18.7	28.8	40.8	53.2	60.2	73.8	
女性	20-29歳	40	22.1	11.6	1.75	1.3	7.3	7.6	16.3	19.3	27.8	37.9	48.0	59.5
	30-39歳	87	27.2	12.6	1.78	6.4	8.6	10.6	16.1	27.9	34.2	42.9	54.2	59.4
	40-49歳	85	26.9	13.6	1.18	0.7	6.7	12.5	16.7	25.2	32.5	47.1	56.3	65.7
	50-59歳	97	25.8	15.1	1.43	1.3	7.6	8.6	13.8	24.1	34.3	46.7	51.1	84.7
	60-69歳	145	26.5	15.0	1.49	2.7	7.4	10.4	15.4	24.2	35.5	48.6	60.8	67.5
	70歳以上	187	23.5	11.6	1.17	1.5	6.0	8.7	15.7	22.8	30.5	40.3	44.3	51.2
	(再掲)20歳以上※	641	25.1	13.2	0.64	1.6	7.3	9.7	15.9	24.1	32.4	43.8	50.1	63.8
	(再掲)65-74歳	151	26.8	14.5	1.45	2.3	7.8	11.3	16.6	23.8	34.7	49.7	59.9	67.1
	(再掲)75歳以上	137	22.9	11.4	1.16	1.2	5.9	8.3	14.6	21.6	30.0	40.0	44.2	50.7
	(再掲)20-49歳※	212	25.4	12.8	0.67	1.1	7.4	9.6	16.6	24.3	32.5	44.1	52.1	62.6
(再掲)40-74歳※	377	26.2	14.2	0.80	2.3	7.5	10.1	16.0	24.4	33.2	47.1	52.0	67.1	
男女計	20-29歳	82	24.8	12.7	1.59	1.3	7.4	8.3	16.3	24.3	32.8	45.7	49.6	59.5
	30-39歳	145	31.6	17.0	1.53	6.9	9.8	11.9	18.7	31.5	37.9	52.1	61.3	106.2
	40-49歳	181	29.1	16.0	1.08	0.2	7.7	10.3	17.5	25.8	36.9	49.1	62.3	87.6
	50-59歳	181	28.6	15.3	1.22	5.2	8.6	10.8	16.0	25.5	38.9	49.1	56.0	75.0
	60-69歳	260	28.4	15.2	1.28	3.2	8.1	10.7	17.2	26.2	36.8	49.1	60.4	68.8
	70歳以上	355	25.3	13.2	1.05	1.6	6.0	9.2	15.9	24.1	33.3	44.3	50.2	60.0
	(再掲)20歳以上※	1,204	27.6	14.9	0.63	2.1	7.6	10.1	16.8	25.5	35.7	47.4	55.6	68.4
	(再掲)65-74歳	281	28.5	14.8	1.32	2.9	8.0	11.1	17.9	26.7	37.2	50.0	59.9	68.0
	(再掲)75歳以上	244	24.7	13.0	1.23	1.3	5.5	8.5	15.1	23.5	32.7	44.1	47.8	59.6
	(再掲)20-49歳※	408	28.4	15.5	0.81	2.0	8.1	9.9	17.7	26.2	35.6	48.7	56.3	73.5
(再掲)40-74歳※	733	28.4	15.2	0.71	2.8	8.1	10.7	16.8	25.7	37.5	48.8	58.6	70.8	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

n-3系脂肪酸(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	2.47	1.47	0.239	0.18	0.44	0.92	1.30	2.25	3.48	4.41	4.79	8.53
	30-39歳	58	2.48	1.39	0.183	0.10	0.65	0.97	1.42	2.10	3.28	4.36	5.24	6.80
	40-49歳	96	2.12	1.29	0.151	0.15	0.48	0.89	1.32	1.84	2.54	4.22	5.58	6.14
	50-59歳	84	2.40	1.24	0.151	0.29	0.69	0.87	1.45	2.16	3.34	4.08	4.79	6.87
	60-69歳	115	2.91	1.72	0.193	0.22	0.91	1.22	1.60	2.52	3.94	5.08	5.87	9.46
	70歳以上	168	2.58	1.68	0.124	0.31	0.72	0.83	1.45	2.28	3.29	4.64	5.71	10.29
	(再掲)20歳以上※	563	2.50	1.51	0.080	0.25	0.72	0.92	1.44	2.19	3.29	4.43	5.28	7.75
	(再掲)65-74歳	130	2.91	1.74	0.205	0.32	0.76	1.01	1.58	2.65	3.89	5.12	6.46	8.86
	(再掲)75歳以上	107	2.55	1.71	0.137	0.58	0.73	0.84	1.47	2.28	3.29	4.11	5.65	10.37
	(再掲)20-49歳※	196	2.34	1.39	0.105	0.14	0.60	0.93	1.32	1.99	3.07	4.37	4.99	7.06
(再掲)40-74歳※	356	2.51	1.50	0.086	0.25	0.73	0.91	1.43	2.13	3.28	4.62	5.54	7.64	
女性	20-29歳	40	1.94	0.96	0.218	0.24	0.36	0.73	1.31	1.79	2.52	3.45	3.70	4.23
	30-39歳	87	1.93	1.13	0.099	0.41	0.55	0.69	1.17	1.58	2.53	3.61	4.04	5.82
	40-49歳	85	1.77	0.88	0.128	0.50	0.75	0.88	1.08	1.59	2.24	3.09	3.63	4.84
	50-59歳	97	2.15	1.04	0.110	0.36	0.78	0.98	1.29	1.95	2.64	3.65	4.36	5.48
	60-69歳	145	2.49	1.44	0.139	0.35	0.65	1.07	1.45	2.28	3.26	4.18	5.11	7.89
	70歳以上	187	2.29	1.43	0.150	0.35	0.69	0.88	1.40	1.96	2.88	4.09	5.36	7.66
	(再掲)20歳以上※	641	2.13	1.23	0.073	0.29	0.69	0.88	1.26	1.84	2.70	3.68	4.34	5.98
	(再掲)65-74歳	151	2.49	1.41	0.105	0.36	0.68	0.96	1.49	2.16	3.29	4.17	5.13	7.65
	(再掲)75歳以上	137	2.28	1.51	0.172	0.14	0.74	0.88	1.30	1.84	2.82	4.12	5.44	8.10
	(再掲)20-49歳※	212	1.87	0.98	0.093	0.26	0.68	0.76	1.17	1.69	2.41	3.43	3.74	4.90
(再掲)40-74歳※	377	2.17	1.19	0.076	0.45	0.75	0.93	1.26	1.95	2.77	3.74	4.38	5.86	
男女計	20-29歳	82	2.21	1.27	0.195	0.18	0.46	0.76	1.32	1.89	3.04	3.66	4.43	8.53
	30-39歳	145	2.21	1.30	0.124	0.33	0.64	0.82	1.23	1.87	3.03	4.00	4.49	6.48
	40-49歳	181	1.95	1.12	0.130	0.29	0.73	0.88	1.19	1.65	2.42	3.30	4.42	5.79
	50-59歳	181	2.28	1.15	0.117	0.30	0.72	0.95	1.43	2.02	3.10	3.82	4.46	5.54
	60-69歳	260	2.70	1.59	0.144	0.26	0.77	1.09	1.52	2.34	3.59	4.81	5.66	8.95
	70歳以上	355	2.41	1.54	0.116	0.38	0.72	0.86	1.43	2.05	3.04	4.22	5.43	8.38
	(再掲)20歳以上※	1,204	2.31	1.39	0.068	0.29	0.69	0.90	1.33	1.99	3.04	4.08	4.84	7.11
	(再掲)65-74歳	281	2.68	1.58	0.137	0.36	0.74	0.99	1.51	2.30	3.54	4.82	5.72	8.12
	(再掲)75歳以上	244	2.38	1.59	0.137	0.41	0.74	0.88	1.41	1.99	3.02	4.12	5.41	9.56
	(再掲)20-49歳※	408	2.11	1.23	0.090	0.24	0.65	0.88	1.24	1.79	2.74	3.73	4.41	6.05
(再掲)40-74歳※	733	2.34	1.37	0.074	0.30	0.74	0.91	1.34	2.02	3.03	4.19	5.00	6.73	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

n-6系脂肪酸(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	11.26	5.27	0.747	3.02	3.69	4.58	6.69	11.66	13.73	18.56	23.14	25.02
	30-39歳	58	11.64	5.33	0.741	1.31	4.68	5.54	7.73	10.78	14.77	19.31	22.77	27.08
	40-49歳	96	11.15	4.02	0.532	2.63	4.76	5.84	8.20	10.56	14.78	16.97	17.66	19.72
	50-59歳	84	11.50	5.03	0.560	3.25	4.78	5.39	8.43	10.87	13.86	19.06	22.89	25.68
	60-69歳	115	10.61	5.54	0.656	2.92	4.40	5.24	7.26	9.48	12.57	16.70	22.22	31.85
	70歳以上	168	9.09	4.56	0.419	2.59	3.19	4.28	5.68	7.89	11.53	16.43	18.06	24.79
	(再掲)20歳以上※	563	10.73	5.01	0.330	2.80	4.15	5.08	7.12	9.93	13.31	17.30	19.10	26.02
	(再掲)65-74歳	130	9.69	4.21	0.345	2.73	4.38	5.02	6.86	9.11	11.98	14.55	17.45	26.10
	(再掲)75歳以上	107	8.65	4.24	0.562	2.12	3.00	4.03	5.43	7.68	10.79	16.39	17.91	19.51
	(再掲)20-49歳※	196	11.33	4.83	0.445	2.54	4.27	5.39	7.77	10.94	14.68	17.46	19.85	24.98
(再掲)40-74歳※	356	10.90	4.93	0.400	2.90	4.66	5.40	7.66	10.09	13.31	16.94	19.07	29.15	
女性	20-29歳	40	9.13	3.19	0.613	2.09	3.17	4.53	6.91	9.98	11.28	13.20	13.76	15.16
	30-39歳	87	9.10	3.82	0.598	1.72	4.31	4.73	6.28	8.54	11.29	14.48	17.35	19.48
	40-49歳	85	9.72	3.77	0.430	3.78	4.56	5.24	6.60	9.15	12.84	15.00	16.94	20.53
	50-59歳	97	10.05	3.95	0.377	2.81	4.66	5.67	7.50	9.51	11.99	14.77	17.17	27.67
	60-69歳	145	8.48	4.57	0.361	0.43	2.76	4.05	5.14	7.32	10.88	15.27	18.57	24.25
	70歳以上	187	7.73	3.49	0.210	1.77	2.30	3.63	5.24	7.15	9.58	12.64	14.38	19.92
	(再掲)20歳以上※	641	8.83	3.88	0.227	1.91	3.44	4.55	6.10	8.36	11.05	13.75	15.91	19.91
	(再掲)65-74歳	151	8.38	4.44	0.311	0.43	2.67	3.76	5.04	7.39	10.63	14.45	17.74	21.97
	(再掲)75歳以上	137	7.44	3.29	0.236	1.65	2.18	3.41	5.16	6.90	9.44	12.28	14.30	17.69
	(再掲)20-49歳※	212	9.34	3.60	0.384	2.16	4.20	5.06	6.65	9.10	11.62	13.77	16.51	19.62
(再掲)40-74歳※	377	9.23	4.15	0.245	1.83	3.81	4.63	6.30	8.62	11.64	14.56	17.16	20.79	
男女計	20-29歳	82	10.23	4.50	0.610	2.18	3.64	4.60	6.86	10.76	12.75	14.91	18.67	25.02
	30-39歳	145	10.39	4.80	0.507	1.53	4.49	4.93	6.83	9.52	13.01	17.33	19.83	25.74
	40-49歳	181	10.45	3.96	0.411	3.59	4.75	5.36	7.71	9.74	13.18	16.83	17.46	19.81
	50-59歳	181	10.78	4.57	0.408	3.21	4.78	5.44	7.88	9.73	12.98	16.05	21.29	25.98
	60-69歳	260	9.53	5.17	0.423	0.62	3.67	4.46	6.15	8.26	11.68	15.72	19.10	30.47
	70歳以上	355	8.30	4.02	0.274	1.99	2.79	3.86	5.50	7.51	10.29	13.72	16.54	19.92
	(再掲)20歳以上※	1,204	9.75	4.56	0.266	2.08	3.71	4.76	6.53	9.10	12.30	15.46	17.83	24.83
	(再掲)65-74歳	281	8.99	4.37	0.294	0.76	3.34	4.39	5.86	8.08	11.35	14.48	17.47	25.00
	(再掲)75歳以上	244	7.91	3.73	0.315	1.82	2.70	3.72	5.32	7.23	9.66	13.03	15.76	18.85
	(再掲)20-49歳※	408	10.36	4.38	0.369	2.41	4.27	5.15	7.08	9.92	12.98	16.78	18.08	24.79
(再掲)40-74歳※	733	10.06	4.62	0.303	2.36	4.20	4.91	6.78	9.34	12.60	16.15	18.19	25.26	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

飽和脂肪酸(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	14.66	6.38	0.777	4.56	5.71	6.65	10.27	13.48	18.78	25.20	29.14	31.75
	30-39歳	58	17.12	7.99	0.877	3.97	6.09	7.73	11.95	15.80	21.95	28.29	33.49	45.52
	40-49歳	96	16.33	7.64	0.600	1.61	4.89	7.49	11.06	15.70	20.33	26.29	28.67	45.42
	50-59歳	84	15.99	6.77	0.749	4.33	6.55	7.92	11.37	14.72	20.31	25.88	28.58	38.37
	60-69歳	115	15.69	7.35	0.863	3.19	5.79	7.85	10.49	14.58	18.98	26.21	31.84	37.84
	70歳以上	168	14.05	7.05	0.707	1.52	3.42	5.29	8.80	13.21	18.82	24.59	26.19	32.32
	(再掲)20歳以上※	563	15.51	7.24	0.296	3.11	5.52	7.38	10.44	14.34	19.55	25.42	28.87	36.42
	(再掲)65-74歳	130	15.17	6.94	0.658	2.73	4.91	5.89	10.34	14.49	19.25	24.10	29.37	34.56
	(再掲)75歳以上	107	13.86	7.13	0.953	1.49	3.09	5.32	8.30	13.20	18.43	24.91	26.22	33.23
	(再掲)20-49歳※	196	15.99	7.37	0.434	3.96	5.83	7.46	10.71	14.46	19.85	25.54	29.65	43.53
(再掲)40-74歳※	356	15.78	7.24	0.371	3.18	5.54	7.73	10.53	14.86	20.18	25.40	28.85	38.09	
女性	20-29歳	40	12.49	5.13	0.881	2.97	5.54	8.49	9.08	10.60	15.90	19.70	22.88	31.47
	30-39歳	87	14.32	6.04	0.972	2.25	5.68	7.10	9.52	13.79	18.14	23.29	25.83	29.60
	40-49歳	85	15.45	7.14	0.695	2.36	6.01	7.34	9.57	14.24	19.31	27.24	30.61	34.45
	50-59歳	97	14.58	6.34	0.621	3.74	5.86	7.57	9.54	13.57	18.17	24.87	27.33	31.95
	60-69歳	145	13.52	7.69	0.755	1.71	4.39	5.51	8.42	11.93	17.16	21.97	28.01	41.90
	70歳以上	187	12.17	5.68	0.470	1.65	4.12	5.90	7.71	11.69	15.28	19.27	22.09	33.89
	(再掲)20歳以上※	641	13.53	6.45	0.284	2.41	5.13	6.37	8.98	12.25	17.29	21.19	25.55	33.76
	(再掲)65-74歳	151	13.45	7.45	0.648	1.91	4.59	6.16	8.07	12.03	16.97	20.64	28.61	41.59
	(再掲)75歳以上	137	11.83	5.73	0.550	1.21	3.67	5.27	7.02	11.58	14.72	19.24	23.00	30.48
	(再掲)20-49歳※	212	14.14	6.31	0.383	2.44	5.87	7.72	9.34	12.98	18.07	22.85	27.83	32.66
(再掲)40-74歳※	377	14.28	6.95	0.388	2.46	5.45	6.78	9.06	13.17	18.14	23.47	27.90	34.90	
男女計	20-29歳	82	13.62	5.88	0.609	2.97	5.83	7.36	9.72	12.83	16.44	21.89	26.18	31.75
	30-39歳	145	15.75	7.21	0.674	3.39	5.90	7.65	10.75	14.70	19.32	25.39	29.37	40.88
	40-49歳	181	15.90	7.39	0.488	2.21	5.52	7.48	10.55	15.18	19.78	26.50	29.55	40.42
	50-59歳	181	15.29	6.58	0.541	4.19	6.37	7.91	10.50	14.08	19.49	25.56	27.63	32.98
	60-69歳	260	14.59	7.59	0.590	2.49	4.89	6.17	9.18	13.80	17.73	24.13	31.38	39.83
	70歳以上	355	12.95	6.35	0.480	1.54	3.90	5.52	8.20	12.17	16.90	21.47	24.75	33.00
	(再掲)20歳以上※	1,204	14.48	6.91	0.253	2.70	5.30	6.76	9.47	13.52	18.35	23.83	27.90	34.54
	(再掲)65-74歳	281	14.25	7.25	0.552	2.43	4.78	6.14	8.84	13.67	17.86	22.24	29.22	36.88
	(再掲)75歳以上	244	12.62	6.38	0.638	1.37	3.56	5.29	7.86	11.87	16.35	21.73	24.85	32.19
	(再掲)20-49歳※	408	15.09	6.93	0.328	2.99	5.89	7.49	9.88	13.77	19.03	24.46	28.46	34.44
(再掲)40-74歳※	733	15.03	7.13	0.335	3.15	5.49	6.98	9.62	14.06	19.15	24.36	28.31	35.64	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

一価不飽和脂肪酸(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	21.16	8.54	0.958	5.09	7.03	9.86	14.44	20.95	27.21	30.84	38.49	42.07
	30-39歳	58	22.65	11.14	1.448	4.43	8.77	9.76	13.80	20.97	29.81	38.60	39.85	70.63
	40-49歳	96	21.17	8.78	0.851	2.55	8.56	10.28	14.74	20.75	27.13	32.90	35.94	51.55
	50-59歳	84	21.74	8.65	0.855	7.29	9.36	10.73	14.83	21.49	27.45	31.31	38.34	54.62
	60-69歳	115	20.27	9.05	1.100	3.90	9.03	10.83	13.17	18.62	25.09	33.20	40.53	49.55
	70歳以上	168	17.77	9.04	0.838	2.01	4.95	7.98	11.44	16.28	22.28	30.10	34.91	48.27
	(再掲)20歳以上※	563	20.54	9.27	0.464	3.65	8.11	9.57	14.06	19.19	26.34	32.16	38.32	46.33
	(再掲)65-74歳	130	19.31	8.36	0.834	2.93	7.78	9.27	13.00	18.64	24.89	31.63	36.01	42.02
	(再掲)75歳以上	107	17.27	8.99	0.975	1.82	4.34	6.81	11.61	16.06	21.22	28.47	34.77	54.44
	(再掲)20-49歳※	196	21.59	9.41	0.638	4.78	8.33	10.10	14.56	20.82	27.57	32.71	38.64	49.45
(再掲)40-74歳※	356	20.69	8.90	0.475	4.06	8.49	10.45	14.06	19.74	25.75	32.82	37.65	45.01	
女性	20-29歳	40	18.33	6.01	0.775	4.38	7.37	9.65	15.21	18.41	22.45	25.67	29.69	36.43
	30-39歳	87	17.51	6.74	1.043	3.20	7.45	8.73	12.81	16.58	21.87	26.27	30.01	35.70
	40-49歳	85	19.13	7.58	0.770	3.62	8.77	10.09	12.86	18.58	25.63	29.09	32.31	36.95
	50-59歳	97	19.07	8.24	0.732	6.03	7.90	10.22	13.67	17.94	22.89	28.49	39.49	45.38
	60-69歳	145	17.16	10.47	1.049	1.92	5.72	8.05	10.48	15.07	20.89	28.91	33.40	62.68
	70歳以上	187	15.00	6.66	0.553	2.15	5.67	7.01	9.89	14.26	18.59	23.58	27.59	33.91
	(再掲)20歳以上※	641	17.30	7.86	0.374	3.32	6.54	8.65	11.97	16.58	21.58	27.44	30.42	43.93
	(再掲)65-74歳	151	17.13	10.01	0.886	2.40	6.59	8.40	10.59	15.40	20.76	29.15	33.88	61.47
	(再掲)75歳以上	137	14.44	6.61	0.601	1.48	4.72	6.45	9.12	13.80	18.56	23.20	27.25	32.68
	(再掲)20-49歳※	212	18.40	6.85	0.498	3.83	8.04	9.74	13.20	17.80	22.81	27.71	30.50	36.30
(再掲)40-74歳※	377	18.10	8.67	0.484	3.97	7.46	9.06	12.16	16.49	22.79	28.32	32.31	45.28	
男女計	20-29歳	82	19.80	7.53	0.724	4.38	7.79	9.84	14.69	18.61	24.61	29.55	35.97	42.07
	30-39歳	145	20.13	9.56	0.986	3.96	8.36	9.29	12.93	18.93	26.25	31.31	38.49	58.99
	40-49歳	181	20.17	8.26	0.698	3.86	8.69	10.39	13.35	19.89	26.02	30.85	33.45	39.93
	50-59歳	181	20.41	8.53	0.571	6.51	9.05	10.55	14.10	19.30	24.76	30.23	38.31	45.36
	60-69歳	260	18.69	9.91	0.828	3.34	7.46	8.92	11.97	16.61	22.82	31.47	36.84	56.28
	70歳以上	355	16.15	7.85	0.558	2.08	5.49	7.18	10.59	15.20	19.86	26.96	31.59	38.38
	(再掲)20歳以上※	1,204	18.86	8.72	0.389	3.45	7.10	8.97	12.73	17.72	23.69	29.88	34.12	44.57
	(再掲)65-74歳	281	18.15	9.33	0.730	3.23	7.03	8.52	11.96	16.48	22.38	29.96	34.48	48.64
	(再掲)75歳以上	244	15.54	7.73	0.654	1.72	4.56	6.60	9.84	14.97	19.12	25.30	29.89	35.76
	(再掲)20-49歳※	408	20.03	8.40	0.503	4.42	8.20	9.96	13.79	19.46	25.67	30.67	34.18	40.97
(再掲)40-74歳※	733	19.39	8.88	0.410	4.24	8.04	9.61	12.86	18.26	24.29	30.65	34.50	45.13	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

多価不飽和脂肪酸(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	13.89	6.08	0.864	3.21	4.43	5.70	9.15	14.21	17.51	22.32	26.84	29.43
	30-39歳	58	14.25	6.04	0.852	1.41	5.54	7.36	10.02	13.53	18.42	22.70	26.65	32.03
	40-49歳	96	13.60	4.92	0.657	2.90	5.88	7.67	10.06	12.84	18.00	20.49	22.14	23.71
	50-59歳	84	14.20	6.00	0.643	3.70	5.59	6.28	10.47	13.79	16.60	23.67	27.06	35.45
	60-69歳	115	13.73	6.29	0.713	3.35	6.31	7.71	9.65	12.66	15.39	21.24	27.34	37.24
	70歳以上	168	11.89	5.57	0.503	2.95	4.82	5.87	8.01	10.67	14.83	19.52	22.23	30.85
	(再掲)20歳以上※	563	13.46	5.84	0.372	3.23	5.34	6.72	9.35	12.83	16.51	21.04	23.81	31.96
	(再掲)65-74歳	130	12.83	4.98	0.444	3.25	6.08	7.19	9.39	12.55	14.95	18.74	21.38	32.22
	(再掲)75歳以上	107	11.42	5.24	0.640	3.24	4.72	5.36	7.58	10.44	13.73	19.27	21.84	28.70
	(再掲)20-49歳※	196	13.88	5.63	0.518	2.71	5.40	7.16	9.76	13.56	18.09	21.16	23.77	29.06
(再掲)40-74歳※	356	13.67	5.80	0.426	3.63	5.84	7.34	9.81	12.97	16.67	20.84	23.70	35.68	
女性	20-29歳	40	11.14	3.81	0.798	2.33	4.59	6.29	7.76	11.66	14.31	16.03	16.65	18.74
	30-39歳	87	11.20	4.33	0.670	2.54	5.41	6.04	8.09	10.84	13.47	17.70	20.98	23.07
	40-49歳	85	11.77	4.32	0.519	4.67	5.82	6.54	9.02	11.28	15.07	18.18	20.11	24.63
	50-59歳	97	12.35	4.49	0.425	3.17	6.19	7.15	9.31	11.71	14.52	17.55	19.70	29.33
	60-69歳	145	11.17	5.21	0.427	1.03	4.11	6.14	7.88	9.95	13.48	18.49	21.88	30.17
	70歳以上	187	10.25	4.31	0.294	2.16	4.01	5.21	7.49	9.80	12.92	16.43	17.42	25.57
	(再掲)20歳以上※	641	11.16	4.48	0.271	2.59	4.95	6.00	7.95	10.61	13.78	17.00	18.88	24.76
	(再掲)65-74歳	151	11.03	5.13	0.380	1.20	4.22	5.80	7.87	10.39	13.41	17.67	21.44	28.15
	(再掲)75歳以上	137	9.99	4.18	0.338	1.78	3.36	5.05	7.49	9.43	12.69	16.39	17.02	22.27
	(再掲)20-49歳※	212	11.40	4.15	0.444	2.79	5.49	6.30	8.35	11.00	14.14	16.97	18.81	23.08
(再掲)40-74歳※	377	11.60	4.71	0.270	3.12	5.61	6.32	8.35	10.80	14.19	17.70	20.37	26.23	
男女計	20-29歳	82	12.57	5.27	0.743	2.48	4.75	5.94	8.70	12.81	15.80	18.51	22.41	29.43
	30-39歳	145	12.75	5.47	0.590	2.01	5.56	6.41	8.51	11.75	15.85	20.25	23.22	29.20
	40-49歳	181	12.70	4.71	0.516	4.55	5.95	7.21	9.27	11.64	15.57	20.06	21.49	23.86
	50-59歳	181	13.28	5.37	0.474	3.67	5.92	6.90	10.04	12.71	16.20	19.58	24.95	29.50
	60-69歳	260	12.43	5.90	0.463	2.64	5.15	6.54	8.86	11.28	14.52	19.46	22.46	36.13
	70歳以上	355	10.94	4.94	0.335	2.73	4.39	5.40	7.58	10.38	13.51	17.39	19.92	26.90
	(再掲)20歳以上※	1,204	12.27	5.31	0.301	2.96	5.15	6.24	8.59	11.51	15.06	18.95	22.05	29.37
	(再掲)65-74歳	281	11.87	5.13	0.367	2.89	4.83	6.28	8.12	11.21	14.31	18.45	21.38	30.79
	(再掲)75歳以上	244	10.55	4.67	0.382	2.57	4.07	5.17	7.53	9.86	13.14	16.96	19.50	25.07
	(再掲)20-49歳※	408	12.67	5.11	0.431	2.95	5.54	6.57	8.98	12.04	15.74	19.88	22.12	27.20
(再掲)40-74歳※	733	12.63	5.38	0.317	3.41	5.70	6.67	9.06	11.67	15.32	19.58	22.13	30.95	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

飽和脂肪酸エネルギー比率(%)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	6.85	2.52	0.286	3.04	4.07	4.34	5.41	5.77	8.38	11.12	12.57	13.43
	30-39歳	58	6.80	2.00	0.198	3.12	3.61	4.20	5.28	6.77	8.04	9.29	10.08	12.57
	40-49歳	96	7.02	2.66	0.233	1.04	3.05	3.89	5.17	6.53	9.16	10.35	11.46	17.28
	50-59歳	84	6.64	2.15	0.283	2.58	2.98	3.96	5.26	6.35	8.03	9.48	11.09	12.61
	60-69歳	115	6.27	2.44	0.306	1.85	3.07	3.38	4.52	5.96	7.37	9.70	10.72	14.85
	70歳以上	168	5.97	2.50	0.237	1.12	2.20	2.76	4.24	5.74	7.46	9.83	10.55	10.97
	(再掲)20歳以上※	563	6.55	2.43	0.071	1.70	3.04	3.81	4.84	6.19	8.07	9.81	10.92	13.34
	(再掲)65-74歳	130	6.10	2.34	0.198	1.55	2.41	3.11	4.43	5.87	7.61	9.36	10.52	11.88
	(再掲)75歳以上	107	6.04	2.60	0.326	1.01	1.86	2.80	4.24	5.82	7.49	10.28	10.57	13.28
	(再掲)20-49歳※	196	6.90	2.43	0.128	2.26	3.56	4.29	5.32	6.38	8.54	10.12	11.47	13.73
(再掲)40-74歳※	356	6.53	2.45	0.115	1.84	2.91	3.57	4.88	6.20	8.09	9.58	10.87	13.71	
女性	20-29歳	40	7.38	2.30	0.410	1.86	4.34	4.93	5.63	7.26	8.95	10.42	11.89	12.51
	30-39歳	87	7.66	2.43	0.358	2.04	3.71	4.89	6.05	7.17	9.04	10.96	12.10	16.15
	40-49歳	85	7.65	2.68	0.223	1.61	3.69	4.82	5.66	6.99	9.58	11.14	12.07	14.59
	50-59歳	97	7.53	2.82	0.277	2.75	3.67	4.26	5.25	7.01	9.09	11.54	14.09	15.04
	60-69歳	145	6.69	2.66	0.262	1.99	2.75	3.32	4.58	6.73	8.33	10.41	12.19	13.38
	70歳以上	187	6.58	2.43	0.221	1.37	2.98	3.64	4.78	6.40	8.33	9.79	10.87	12.85
	(再掲)20歳以上※	641	7.13	2.58	0.112	1.83	3.37	3.99	5.22	6.87	8.78	10.63	11.84	14.33
	(再掲)65-74歳	151	6.71	2.57	0.224	2.46	2.96	3.56	4.74	6.68	8.27	10.19	11.77	14.36
	(再掲)75歳以上	137	6.50	2.49	0.288	1.26	2.68	3.56	4.55	6.12	8.39	9.80	10.92	12.55
	(再掲)20-49歳※	212	7.56	2.48	0.150	1.74	3.71	4.88	5.72	7.19	9.26	10.93	11.92	14.13
(再掲)40-74歳※	377	7.20	2.68	0.143	2.47	3.33	3.98	5.18	6.91	8.84	10.82	12.07	14.40	
男女計	20-29歳	82	7.11	2.42	0.266	1.86	4.12	4.43	5.48	6.27	8.64	10.78	12.00	13.43
	30-39歳	145	7.22	2.26	0.208	2.72	3.73	4.51	5.56	7.05	8.50	10.10	11.72	13.48
	40-49歳	181	7.33	2.68	0.157	1.52	3.31	4.02	5.48	6.86	9.24	10.83	11.81	14.41
	50-59歳	181	7.09	2.54	0.266	2.60	3.51	4.13	5.26	6.66	8.44	10.69	11.87	14.54
	60-69歳	260	6.49	2.56	0.197	1.98	2.99	3.33	4.57	6.14	8.06	10.08	11.69	13.80
	70歳以上	355	6.33	2.47	0.201	1.31	2.49	3.53	4.44	6.08	8.00	9.79	10.58	12.48
	(再掲)20歳以上※	1,204	6.85	2.53	0.079	1.83	3.21	3.88	5.12	6.55	8.45	10.30	11.48	14.04
	(再掲)65-74歳	281	6.42	2.48	0.187	1.91	2.63	3.33	4.68	6.06	8.10	9.67	10.70	13.00
	(再掲)75歳以上	244	6.32	2.54	0.277	1.22	2.39	3.50	4.40	6.08	8.06	9.96	10.73	12.45
	(再掲)20-49歳※	408	7.22	2.47	0.105	1.85	3.71	4.37	5.50	6.79	8.96	10.66	11.83	13.91
(再掲)40-74歳※	733	6.87	2.59	0.112	2.06	3.10	3.71	5.04	6.57	8.45	10.35	11.53	14.35	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

P/S比

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	1.05	0.45	0.075	0.24	0.28	0.40	0.71	1.06	1.46	1.68	1.74	1.76
	30-39歳	58	0.90	0.36	0.045	0.21	0.48	0.55	0.64	0.84	1.09	1.36	1.76	2.01
	40-49歳	96	0.98	0.51	0.048	0.31	0.43	0.46	0.64	0.80	1.22	1.59	1.88	3.17
	50-59歳	84	0.95	0.34	0.039	0.22	0.42	0.56	0.71	0.89	1.13	1.47	1.64	2.16
	60-69歳	115	1.01	0.50	0.054	0.24	0.35	0.43	0.66	0.89	1.24	1.66	1.95	2.90
	70歳以上	168	0.99	0.48	0.036	0.33	0.37	0.46	0.64	0.89	1.28	1.69	1.89	2.49
	(再掲)20歳以上※	563	0.98	0.45	0.019	0.26	0.40	0.47	0.67	0.89	1.23	1.62	1.79	2.40
	(再掲)65-74歳	130	1.00	0.50	0.038	0.29	0.37	0.42	0.64	0.89	1.27	1.75	1.95	2.73
	(再掲)75歳以上	107	0.97	0.46	0.043	0.31	0.36	0.44	0.63	0.90	1.24	1.68	1.87	2.37
	(再掲)20-49歳※	196	0.98	0.45	0.038	0.23	0.41	0.47	0.67	0.87	1.23	1.61	1.75	2.60
(再掲)40-74歳※	356	0.99	0.47	0.022	0.27	0.41	0.49	0.66	0.89	1.22	1.61	1.84	2.74	
女性	20-29歳	40	0.98	0.43	0.084	0.29	0.43	0.51	0.67	0.90	1.33	1.49	1.60	2.56
	30-39歳	87	0.88	0.40	0.053	0.32	0.34	0.40	0.58	0.82	1.02	1.39	1.82	2.17
	40-49歳	85	0.86	0.34	0.033	0.25	0.40	0.47	0.66	0.81	1.04	1.33	1.54	2.08
	50-59歳	97	0.95	0.42	0.048	0.31	0.43	0.50	0.66	0.87	1.13	1.49	1.66	3.17
	60-69歳	145	0.96	0.44	0.044	0.22	0.36	0.46	0.65	0.89	1.24	1.58	1.79	2.37
	70歳以上	187	0.95	0.43	0.035	0.33	0.43	0.49	0.60	0.86	1.18	1.54	1.80	2.40
	(再掲)20歳以上※	641	0.93	0.42	0.021	0.29	0.40	0.49	0.63	0.86	1.15	1.47	1.68	2.37
	(再掲)65-74歳	151	0.94	0.42	0.032	0.24	0.37	0.46	0.63	0.89	1.18	1.59	1.79	2.09
	(再掲)75歳以上	137	0.96	0.44	0.046	0.34	0.46	0.50	0.60	0.86	1.18	1.51	1.82	2.46
	(再掲)20-49歳※	212	0.91	0.39	0.030	0.29	0.40	0.48	0.64	0.86	1.11	1.39	1.59	2.29
(再掲)40-74歳※	377	0.92	0.41	0.020	0.29	0.40	0.47	0.64	0.85	1.15	1.47	1.68	2.12	
男女計	20-29歳	82	1.02	0.44	0.068	0.24	0.37	0.47	0.68	1.00	1.37	1.60	1.73	2.56
	30-39歳	145	0.89	0.38	0.036	0.30	0.38	0.49	0.60	0.84	1.05	1.37	1.76	2.04
	40-49歳	181	0.92	0.44	0.036	0.26	0.41	0.47	0.65	0.81	1.15	1.50	1.75	2.77
	50-59歳	181	0.95	0.38	0.038	0.29	0.44	0.52	0.69	0.89	1.13	1.47	1.66	2.11
	60-69歳	260	0.98	0.47	0.040	0.25	0.36	0.45	0.65	0.89	1.24	1.63	1.91	2.42
	70歳以上	355	0.97	0.45	0.025	0.33	0.41	0.49	0.63	0.87	1.20	1.64	1.87	2.39
	(再掲)20歳以上※	1,204	0.96	0.43	0.018	0.29	0.40	0.49	0.65	0.87	1.18	1.54	1.75	2.37
	(再掲)65-74歳	281	0.96	0.45	0.031	0.28	0.37	0.45	0.63	0.89	1.19	1.64	1.87	2.32
	(再掲)75歳以上	244	0.96	0.45	0.035	0.33	0.42	0.49	0.62	0.87	1.19	1.62	1.85	2.40
	(再掲)20-49歳※	408	0.95	0.42	0.031	0.26	0.40	0.48	0.66	0.87	1.16	1.53	1.72	2.28
(再掲)40-74歳※	733	0.96	0.44	0.019	0.29	0.40	0.47	0.66	0.87	1.17	1.55	1.77	2.39	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

コレステロール(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	340	176	22.5	46	56	104	222	326	435	604	669	904
	30-39歳	58	365	192	28.2	59	83	101	220	369	484	597	667	1,067
	40-49歳	96	312	176	15.1	11	55	101	163	279	436	569	648	779
	50-59歳	84	341	184	17.6	61	92	123	226	340	412	550	621	1,216
	60-69歳	115	314	177	15.8	40	77	106	174	283	421	563	665	885
	70歳以上	168	337	174	12.1	24	68	118	194	329	457	555	603	847
	(再掲)20歳以上※	563	333	179	8.1	32	75	109	193	326	442	559	645	897
	(再掲)65-74歳	130	336	182	13.8	43	67	122	197	314	430	565	669	1,001
	(再掲)75歳以上	107	333	170	14.8	19	71	118	188	316	478	561	601	739
	(再掲)20-49歳※	196	336	181	13.0	31	70	102	198	329	447	581	657	912
(再掲)40-74歳※	356	324	179	7.4	31	80	109	180	313	423	554	638	872	
女性	20-29歳	40	302	168	28.5	17	42	77	203	320	378	477	537	1,121
	30-39歳	87	281	154	16.7	41	66	80	171	273	352	505	539	879
	40-49歳	85	273	149	12.3	11	78	106	157	257	350	449	593	722
	50-59歳	97	315	175	17.2	28	81	109	172	322	427	491	675	958
	60-69歳	145	279	180	12.7	38	56	88	132	249	378	526	601	948
	70歳以上	187	287	159	9.8	33	55	85	158	286	399	504	543	755
	(再掲)20歳以上※	641	289	164	7.4	21	63	91	162	287	381	496	549	819
	(再掲)65-74歳	151	297	183	15.8	22	64	88	151	289	395	539	606	928
	(再掲)75歳以上	137	286	153	12.3	35	55	87	162	286	388	501	546	697
	(再掲)20-49歳※	212	285	156	10.3	16	69	91	169	291	362	471	535	826
(再掲)40-74歳※	377	288	170	7.9	24	72	99	151	276	381	498	574	874	
男女計	20-29歳	82	322	172	19.8	17	57	82	217	324	400	507	658	1,121
	30-39歳	145	324	179	18.4	42	75	96	182	316	443	530	608	983
	40-49歳	181	293	164	10.3	14	71	104	160	262	379	508	638	766
	50-59歳	181	328	179	14.9	28	84	114	176	334	423	538	619	1,113
	60-69歳	260	296	179	11.0	40	62	97	162	274	399	541	637	869
	70歳以上	355	308	167	8.5	25	62	95	181	297	418	538	589	757
	(再掲)20歳以上※	1,204	310	173	6.9	29	70	101	174	304	411	525	611	857
	(再掲)65-74歳	281	316	183	12.2	28	68	100	174	296	411	546	654	891
	(再掲)75歳以上	244	304	161	10.0	33	61	96	181	293	417	525	590	738
	(再掲)20-49歳※	408	311	171	9.9	20	70	99	175	311	410	516	628	880
(再掲)40-74歳※	733	306	175	6.6	28	72	102	167	292	407	536	629	875	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

炭水化物(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	267.4	76.7	8.61	128.3	151.4	176.9	214.5	250.3	304.8	387.3	440.3	475.3
	30-39歳	58	309.8	103.6	12.32	103.6	176.0	184.8	217.3	296.6	384.3	459.4	501.7	640.4
	40-49歳	96	279.5	80.8	9.21	90.9	170.0	182.0	220.7	269.9	337.2	392.5	421.3	510.5
	50-59歳	84	290.1	71.4	8.95	133.0	186.6	219.7	246.1	277.4	326.3	382.5	433.7	536.2
	60-69歳	115	303.7	69.7	8.05	145.8	181.9	218.9	263.0	296.1	339.3	402.3	432.2	502.2
	70歳以上	168	300.9	84.7	8.28	117.7	185.2	198.1	240.5	294.7	348.2	420.1	463.3	547.8
	(再掲)20歳以上※	563	292.1	82.3	4.45	124.5	174.5	197.5	236.8	278.8	340.6	401.3	447.5	513.9
	(再掲)65-74歳	130	308.7	72.5	8.67	161.1	182.0	224.7	266.4	303.4	344.2	405.3	439.7	519.0
	(再掲)75歳以上	107	295.1	84.6	9.90	122.6	186.0	197.5	235.2	283.6	345.5	414.5	462.1	571.0
	(再掲)20-49歳※	196	284.0	87.9	6.38	103.7	163.9	185.3	216.7	268.4	340.0	398.8	444.6	554.4
(再掲)40-74歳※	356	294.0	76.4	5.21	117.6	174.6	203.2	246.4	286.0	339.9	398.5	433.1	496.2	
女性	20-29歳	40	208.2	68.0	15.47	77.0	83.0	92.8	151.7	213.4	255.7	287.0	321.3	342.0
	30-39歳	87	230.2	59.6	6.82	79.7	144.3	157.5	189.8	231.5	260.0	307.0	323.2	501.6
	40-49歳	85	242.6	59.6	8.08	119.8	146.6	166.6	195.3	240.4	289.0	318.5	331.2	433.8
	50-59歳	97	230.9	57.8	5.73	40.2	136.5	154.1	201.9	231.4	268.9	303.4	319.7	384.4
	60-69歳	145	249.8	68.8	6.70	77.2	130.9	169.6	213.7	245.5	283.8	348.5	363.1	462.6
	70歳以上	187	240.3	63.9	4.90	128.7	149.8	165.7	194.1	235.5	276.5	328.0	356.8	444.6
	(再掲)20歳以上※	641	235.4	64.4	3.89	84.0	135.1	154.7	195.3	235.7	273.7	315.3	344.4	402.8
	(再掲)65-74歳	151	249.2	64.8	5.86	84.4	153.4	183.6	209.2	242.4	286.1	339.1	362.7	463.5
	(再掲)75歳以上	137	237.3	66.0	6.73	122.3	138.4	157.4	184.7	236.0	273.8	331.0	357.1	442.9
	(再掲)20-49歳※	212	227.6	63.9	7.01	77.4	104.9	148.1	187.1	230.1	271.3	311.1	326.8	386.0
(再掲)40-74歳※	377	242.7	62.2	3.51	90.1	148.0	172.1	204.9	239.6	281.9	319.0	357.0	434.9	
男女計	20-29歳	82	238.9	78.1	10.04	77.2	93.3	145.8	204.4	241.2	275.6	346.0	399.6	475.3
	30-39歳	145	270.8	93.5	8.50	99.9	152.9	170.6	207.7	254.1	322.7	392.7	461.1	608.7
	40-49歳	181	261.3	73.4	7.45	103.4	148.3	174.7	207.5	255.8	303.5	366.1	395.2	469.6
	50-59歳	181	260.5	71.2	7.24	77.1	150.4	177.9	218.2	253.6	297.6	354.9	387.3	478.6
	60-69歳	260	276.4	74.2	6.37	94.3	161.4	186.8	229.2	272.8	318.3	367.0	408.8	482.2
	70歳以上	355	265.6	79.1	5.81	130.8	157.4	171.1	209.2	254.6	313.2	361.4	410.9	511.1
	(再掲)20歳以上※	1,204	262.8	78.8	3.91	92.7	149.2	171.1	212.4	253.9	306.2	363.9	403.6	490.4
	(再掲)65-74歳	281	277.0	74.6	6.22	102.1	171.1	187.8	225.2	271.4	320.3	372.9	414.4	487.8
	(再掲)75歳以上	244	259.8	78.9	6.90	126.5	149.7	166.4	203.3	250.8	306.8	358.2	407.3	511.7
	(再掲)20-49歳※	408	256.5	82.0	6.15	85.6	137.1	159.2	207.0	247.8	299.4	367.3	400.8	491.9
(再掲)40-74歳※	733	268.2	74.1	3.96	102.8	160.9	182.6	218.2	261.7	309.3	366.3	403.5	480.1	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

炭水化物エネルギー比率(%)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	58.2	7.3	1.01	39.6	46.3	49.7	54.0	57.2	63.0	67.5	73.4	77.2
	30-39歳	58	59.7	7.7	1.08	34.4	46.1	49.1	53.4	61.0	65.2	68.2	70.6	78.1
	40-49歳	96	59.5	8.5	1.00	40.5	43.6	47.6	53.9	59.5	66.7	71.1	72.5	79.2
	50-59歳	84	59.7	7.4	0.85	40.4	48.1	50.5	55.0	59.2	65.5	68.6	72.8	76.9
	60-69歳	115	61.3	7.6	0.93	34.7	49.3	52.8	56.4	61.0	66.3	70.3	74.4	78.4
	70歳以上	168	62.8	8.6	0.74	37.3	47.8	52.1	57.8	62.5	67.9	73.5	77.0	83.6
	(再掲)20歳以上※	563	60.4	8.0	0.50	39.9	46.6	50.0	55.4	60.5	65.9	71.0	73.9	78.4
	(再掲)65-74歳	130	61.6	7.7	0.90	40.6	48.8	52.2	56.4	61.5	66.9	71.1	74.3	80.0
	(再掲)75歳以上	107	62.9	8.7	0.99	36.7	48.3	52.5	57.6	62.8	68.2	74.3	77.4	85.9
	(再掲)20-49歳※	196	59.1	7.9	0.74	39.3	46.4	49.0	54.0	59.4	64.6	69.2	71.8	78.1
(再掲)40-74歳※	356	60.5	8.0	0.50	40.4	47.2	50.0	55.6	60.4	66.3	71.1	73.8	78.4	
女性	20-29歳	40	55.3	8.7	1.55	33.4	40.2	41.0	50.2	54.0	60.3	67.5	70.9	77.1
	30-39歳	87	57.5	7.1	0.91	37.3	46.5	49.3	52.5	57.1	62.0	66.2	69.7	75.5
	40-49歳	85	58.4	7.9	0.80	40.3	47.4	49.3	52.1	57.5	64.5	69.5	71.9	80.1
	50-59歳	97	56.6	8.6	0.79	32.1	40.4	44.1	52.0	56.2	62.6	66.3	70.0	76.3
	60-69歳	145	58.5	8.6	0.85	37.3	41.8	47.3	54.5	59.4	64.4	69.2	72.0	78.9
	70歳以上	187	60.0	8.1	0.74	42.0	46.5	49.4	54.1	59.8	64.6	71.2	73.2	78.9
	(再掲)20歳以上※	641	58.1	8.3	0.51	37.6	43.9	48.3	52.6	58.1	63.7	69.3	71.6	77.4
	(再掲)65-74歳	151	58.7	8.3	0.86	37.8	43.1	49.1	54.1	58.6	63.9	69.6	72.3	79.2
	(再掲)75歳以上	137	60.2	8.3	0.90	42.0	46.4	49.0	53.8	61.2	65.2	71.4	73.7	79.6
	(再掲)20-49歳※	212	57.1	8.1	0.76	37.0	41.9	48.2	51.8	56.8	61.9	68.4	70.3	77.0
(再掲)40-74歳※	377	58.1	8.3	0.48	37.5	43.7	48.2	52.6	57.9	63.9	69.2	71.2	77.3	
男女計	20-29歳	82	56.8	8.1	1.02	35.7	40.7	46.4	51.6	56.8	61.9	66.4	70.6	77.2
	30-39歳	145	58.7	7.4	0.75	36.3	46.5	49.2	52.8	59.6	64.5	66.6	69.9	76.7
	40-49歳	181	59.0	8.2	0.65	40.5	46.8	49.0	53.1	58.3	65.2	70.1	72.1	79.5
	50-59歳	181	58.2	8.1	0.72	32.8	43.0	49.5	52.8	58.4	63.4	68.1	72.0	76.6
	60-69歳	260	59.8	8.2	0.69	36.9	44.3	49.5	54.9	60.0	65.4	69.9	72.8	78.1
	70歳以上	355	61.1	8.4	0.65	41.8	47.1	49.6	55.3	61.2	66.9	71.9	74.4	80.2
	(再掲)20歳以上※	1,204	59.2	8.3	0.47	39.4	46.2	49.4	53.7	59.4	64.9	69.7	72.4	78.1
	(再掲)65-74歳	281	60.0	8.2	0.81	38.6	46.0	50.2	54.8	59.8	66.1	70.1	73.0	78.6
	(再掲)75歳以上	244	61.3	8.5	0.81	41.7	46.8	49.5	55.3	61.3	66.9	72.0	75.9	80.6
	(再掲)20-49歳※	408	58.1	8.0	0.64	38.8	46.1	48.9	52.8	58.2	63.7	68.5	71.2	77.2
(再掲)40-74歳※	733	59.3	8.2	0.41	40.0	46.0	49.5	54.0	59.3	65.4	69.8	72.3	78.1	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

総食物繊維(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	12.8	6.0	1.00	3.7	4.4	6.0	8.1	12.9	15.4	18.9	28.9	31.8
	30-39歳	58	15.8	7.2	0.83	3.7	4.4	7.6	11.3	14.5	20.3	25.5	31.4	37.0
	40-49歳	96	14.2	5.2	0.65	4.2	6.9	8.0	10.2	13.5	17.1	20.7	26.1	29.3
	50-59歳	84	16.0	6.3	0.74	2.9	9.1	9.7	11.1	14.9	19.0	26.3	28.5	35.1
	60-69歳	115	18.2	7.4	0.73	4.5	8.8	9.7	13.0	17.3	22.6	27.0	29.3	51.5
	70歳以上	168	19.1	8.3	0.77	4.5	9.1	10.1	12.8	17.4	24.0	31.1	36.8	40.5
	(再掲)20歳以上※	563	16.2	7.2	0.46	4.2	6.5	8.6	11.5	14.6	20.1	26.6	29.8	38.7
	(再掲)65-74歳	130	19.9	7.9	0.76	8.1	9.7	10.8	13.5	18.9	25.1	29.1	34.0	51.8
	(再掲)75歳以上	107	18.1	8.1	1.08	4.2	7.8	9.8	12.2	16.8	21.4	30.9	35.9	40.4
	(再掲)20-49歳※	196	14.2	6.2	0.54	3.7	4.9	6.8	10.4	13.0	17.1	22.0	26.3	32.5
(再掲)40-74歳※	356	16.8	7.0	0.44	5.1	8.0	9.5	11.7	15.4	20.8	26.8	29.1	38.0	
女性	20-29歳	40	12.3	5.7	1.03	2.3	3.5	6.0	9.5	10.7	16.1	19.4	22.9	35.2
	30-39歳	87	14.1	5.3	0.67	5.4	7.1	8.4	10.3	13.3	16.3	20.5	25.0	32.9
	40-49歳	85	13.4	5.7	0.65	4.6	6.5	7.9	9.4	11.6	16.8	20.9	25.8	31.4
	50-59歳	97	15.6	5.4	0.56	5.1	6.8	8.2	12.2	15.2	19.4	23.4	25.1	31.6
	60-69歳	145	19.7	12.0	1.24	1.5	8.0	10.0	13.4	18.0	23.0	28.8	30.5	79.3
	70歳以上	187	16.2	6.7	0.53	5.0	7.2	9.0	11.4	15.5	19.7	24.3	28.0	38.4
	(再掲)20歳以上※	641	15.5	7.7	0.44	4.1	6.8	8.3	10.4	14.4	18.8	24.2	27.7	37.0
	(再掲)65-74歳	151	19.1	11.2	1.39	2.4	7.8	9.5	12.6	17.5	23.3	27.9	29.8	76.1
	(再掲)75歳以上	137	16.1	6.9	0.65	4.9	6.7	8.9	11.6	15.0	18.9	24.5	30.4	41.3
	(再掲)20-49歳※	212	13.3	5.6	0.54	3.2	6.0	6.9	9.5	11.9	16.3	20.0	23.6	33.0
(再掲)40-74歳※	377	16.4	8.5	0.57	4.9	7.2	8.8	10.8	15.3	20.0	25.4	28.6	46.2	
男女計	20-29歳	82	12.6	5.8	0.71	2.6	4.4	6.1	8.9	12.3	15.9	19.0	23.0	35.2
	30-39歳	145	14.9	6.4	0.57	4.2	6.6	8.3	10.8	13.9	18.1	22.8	29.1	35.0
	40-49歳	181	13.8	5.4	0.61	4.6	6.9	8.0	9.7	12.4	17.1	20.7	26.1	29.5
	50-59歳	181	15.8	5.9	0.51	5.1	7.7	9.3	11.8	14.9	19.2	24.3	27.0	33.2
	60-69歳	260	19.0	10.0	0.65	4.4	8.7	10.0	13.1	17.7	22.8	27.8	29.6	67.3
	70歳以上	355	17.4	7.5	0.59	4.9	7.8	9.5	11.9	16.1	21.4	27.7	32.3	40.4
	(再掲)20歳以上※	1,204	15.9	7.5	0.40	4.3	6.7	8.4	10.9	14.5	19.4	25.3	28.9	37.6
	(再掲)65-74歳	281	19.4	9.8	0.79	5.7	9.0	10.0	13.1	18.1	24.1	28.3	30.4	63.9
	(再掲)75歳以上	244	16.8	7.5	0.69	4.8	7.2	9.2	11.8	15.7	20.1	26.7	32.4	40.3
	(再掲)20-49歳※	408	13.7	5.9	0.46	3.6	5.9	6.9	9.8	12.6	16.7	21.4	26.0	32.4
(再掲)40-74歳※	733	16.6	7.8	0.39	5.1	7.5	9.0	11.2	15.4	20.3	26.3	28.8	37.7	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

水溶性食物繊維(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	3.0	1.6	0.26	0.5	0.6	1.2	1.8	2.7	3.6	5.5	7.0	8.0
	30-39歳	58	3.6	1.6	0.19	0.7	1.2	1.4	2.4	3.3	4.7	6.1	6.7	7.4
	40-49歳	96	3.3	1.4	0.17	0.9	1.4	1.6	2.1	3.3	4.2	5.4	5.7	6.5
	50-59歳	84	3.6	1.5	0.21	0.5	1.7	2.1	2.3	3.3	4.8	5.9	6.5	8.5
	60-69歳	115	4.0	1.8	0.18	1.0	1.6	2.0	2.6	3.6	5.1	6.7	7.1	10.6
	70歳以上	168	4.2	2.0	0.17	1.0	1.8	2.1	2.7	3.9	5.3	7.1	8.2	10.9
	(再掲)20歳以上※	563	3.7	1.8	0.11	0.6	1.3	1.7	2.4	3.4	4.8	6.1	6.9	8.6
	(再掲)65-74歳	130	4.4	2.0	0.22	1.6	2.0	2.4	2.8	4.2	5.7	6.8	8.5	12.3
	(再掲)75歳以上	107	3.9	1.9	0.20	0.9	1.7	2.0	2.6	3.6	4.7	7.1	8.1	9.2
	(再掲)20-49歳※	196	3.3	1.6	0.14	0.6	1.2	1.4	2.2	3.1	4.2	5.5	6.2	7.8
(再掲)40-74歳※	356	3.8	1.8	0.12	1.0	1.6	1.9	2.5	3.5	4.9	6.1	6.8	9.1	
女性	20-29歳	40	2.8	1.3	0.24	0.3	1.0	1.2	2.1	2.5	3.5	4.7	5.4	7.4
	30-39歳	87	3.2	1.5	0.18	0.8	1.5	1.8	2.3	2.9	3.9	4.5	6.2	11.4
	40-49歳	85	3.1	1.4	0.18	1.0	1.3	1.6	2.0	2.7	4.1	5.2	5.7	7.3
	50-59歳	97	3.5	1.3	0.14	0.8	1.2	1.7	2.7	3.6	4.2	5.4	6.1	6.6
	60-69歳	145	4.2	1.8	0.14	0.2	1.6	2.2	2.7	4.1	5.5	6.3	7.5	9.7
	70歳以上	187	3.7	1.7	0.12	0.9	1.5	1.8	2.5	3.5	4.6	5.5	6.4	8.7
	(再掲)20歳以上※	641	3.5	1.6	0.10	0.8	1.3	1.7	2.3	3.2	4.4	5.5	6.3	8.4
	(再掲)65-74歳	151	4.0	1.7	0.16	0.5	1.6	1.9	2.5	3.9	5.2	6.3	7.1	8.6
	(再掲)75歳以上	137	3.7	1.8	0.16	0.9	1.3	1.8	2.5	3.5	4.7	5.5	6.3	11.2
	(再掲)20-49歳※	212	3.0	1.4	0.14	0.5	1.3	1.5	2.1	2.7	3.9	4.9	5.4	7.6
(再掲)40-74歳※	377	3.6	1.6	0.08	1.0	1.5	1.8	2.4	3.6	4.6	5.9	6.5	8.4	
男女計	20-29歳	82	2.9	1.5	0.17	0.3	0.7	1.2	2.1	2.7	3.6	5.0	5.6	8.0
	30-39歳	145	3.4	1.6	0.13	0.8	1.2	1.6	2.4	3.0	4.3	5.3	6.5	9.5
	40-49歳	181	3.2	1.4	0.16	1.0	1.3	1.6	2.1	3.0	4.2	5.3	5.7	7.1
	50-59歳	181	3.6	1.4	0.14	0.7	1.6	1.9	2.5	3.4	4.4	5.7	6.3	7.3
	60-69歳	260	4.1	1.8	0.12	1.0	1.6	2.2	2.7	3.9	5.4	6.5	7.2	9.7
	70歳以上	355	3.9	1.8	0.14	0.9	1.6	1.9	2.6	3.6	4.9	6.2	7.2	9.2
	(再掲)20歳以上※	1,204	3.6	1.7	0.09	0.7	1.3	1.7	2.4	3.3	4.5	5.9	6.6	8.5
	(再掲)65-74歳	281	4.2	1.9	0.16	1.1	1.7	2.2	2.7	4.0	5.3	6.5	7.3	10.5
	(再掲)75歳以上	244	3.8	1.8	0.15	0.9	1.5	1.8	2.5	3.5	4.7	5.9	7.2	8.9
	(再掲)20-49歳※	408	3.2	1.5	0.11	0.6	1.2	1.5	2.1	2.9	4.0	5.3	6.0	7.6
(再掲)40-74歳※	733	3.7	1.7	0.09	1.0	1.5	1.8	2.4	3.5	4.8	6.0	6.6	8.6	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

不溶性食物繊維(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	9.4	4.2	0.69	2.7	3.0	3.4	6.4	9.6	10.8	13.9	17.9	22.0
	30-39歳	58	11.5	5.4	0.67	1.9	3.1	5.2	8.2	10.6	15.5	18.1	23.8	30.0
	40-49歳	96	10.3	3.7	0.43	3.5	4.7	5.4	7.6	9.7	12.9	14.7	16.2	23.2
	50-59歳	84	11.8	4.7	0.54	2.6	6.2	6.9	8.3	10.5	13.7	19.9	22.5	27.4
	60-69歳	115	13.3	5.2	0.55	3.0	5.9	7.3	9.7	12.6	17.1	19.6	21.0	35.6
	70歳以上	168	14.0	6.2	0.57	3.4	6.3	7.3	9.6	12.8	17.9	23.4	26.6	30.7
	(再掲)20歳以上※	563	11.9	5.3	0.34	2.8	4.5	6.2	8.3	10.7	14.5	19.0	22.1	27.9
	(再掲)65-74歳	130	14.6	5.7	0.50	6.6	7.3	7.9	10.2	13.7	18.3	21.3	23.8	37.0
	(再掲)75歳以上	107	13.3	6.1	0.77	3.1	5.8	7.1	9.4	11.6	15.8	23.4	26.3	30.5
	(再掲)20-49歳※	196	10.3	4.5	0.38	2.6	3.4	4.6	7.7	9.7	12.7	16.2	18.0	24.9
(再掲)40-74歳※	356	12.3	5.1	0.32	3.6	5.4	6.9	8.7	11.3	14.8	19.3	22.1	27.8	
女性	20-29歳	40	9.2	4.4	0.79	2.0	2.8	3.9	6.8	8.0	12.1	14.3	15.4	27.8
	30-39歳	87	10.3	3.9	0.50	4.1	5.2	6.2	7.6	9.5	12.2	14.6	19.3	23.0
	40-49歳	85	9.8	3.9	0.41	3.4	4.7	5.6	7.0	8.7	12.6	14.6	16.2	22.9
	50-59歳	97	11.6	4.2	0.43	3.8	5.1	6.2	8.6	11.0	13.8	17.7	19.6	23.3
	60-69歳	145	13.6	5.4	0.38	1.4	6.1	7.3	10.0	12.7	17.1	20.9	22.6	32.8
	70歳以上	187	11.9	5.0	0.41	4.0	5.5	6.6	8.4	11.1	14.0	18.6	21.5	29.6
	(再掲)20歳以上※	641	11.3	4.8	0.31	3.1	4.7	6.1	7.6	10.4	13.7	17.9	20.5	26.2
	(再掲)65-74歳	151	13.4	5.2	0.52	1.9	5.9	7.0	9.3	12.9	17.3	20.5	21.9	25.2
	(再掲)75歳以上	137	11.8	5.2	0.49	3.9	4.9	6.6	8.5	10.6	13.8	19.1	22.9	30.0
	(再掲)20-49歳※	212	9.7	4.1	0.39	2.0	4.4	4.9	7.1	9.0	12.2	14.5	16.2	23.2
(再掲)40-74歳※	377	11.8	4.8	0.31	3.6	5.3	6.3	8.0	11.3	14.5	18.4	20.9	25.1	
男女計	20-29歳	82	9.3	4.3	0.50	2.0	3.0	3.9	6.6	9.3	11.6	14.2	16.3	27.8
	30-39歳	145	10.9	4.8	0.44	2.2	4.3	6.2	7.7	10.1	12.5	17.5	19.3	27.6
	40-49歳	181	10.0	3.8	0.39	3.5	4.8	5.6	7.2	9.3	12.6	14.6	16.2	23.0
	50-59歳	181	11.7	4.5	0.36	3.8	5.5	6.8	8.5	10.9	13.6	18.3	20.5	23.8
	60-69歳	260	13.4	5.3	0.35	3.0	6.0	7.3	9.8	12.6	17.1	20.5	21.8	32.5
	70歳以上	355	12.8	5.6	0.46	3.9	5.8	6.9	8.8	11.5	15.7	20.9	24.0	29.6
	(再掲)20歳以上※	1,204	11.6	5.1	0.29	2.9	4.6	6.1	8.0	10.6	13.9	18.4	21.3	27.7
	(再掲)65-74歳	281	14.0	5.5	0.40	4.2	6.6	7.4	10.0	13.4	17.8	20.8	23.2	31.5
	(再掲)75歳以上	244	12.3	5.6	0.52	3.7	5.2	6.7	8.7	11.1	14.0	20.5	24.4	29.9
	(再掲)20-49歳※	408	10.0	4.3	0.32	2.6	4.1	4.9	7.3	9.4	12.3	15.4	17.5	23.9
(再掲)40-74歳※	733	12.1	5.0	0.27	3.7	5.4	6.6	8.3	11.3	14.6	18.8	21.3	26.4	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンA( $\mu$ gRE)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	488	704	91	105	117	130	175	287	532	1,026	1,287	4,904
	30-39歳	58	475	317	47	87	155	180	231	398	572	964	1,168	1,540
	40-49歳	96	435	296	38	42	66	122	243	356	592	839	1,039	1,356
	50-59歳	84	420	278	32	56	123	154	261	386	492	642	949	1,795
	60-69歳	115	528	345	33	57	128	172	291	428	683	1,081	1,289	1,797
	70歳以上	168	604	411	33	81	173	201	318	520	715	1,082	1,548	2,086
	(再掲)20歳以上※	563	500	421	20	59	124	164	252	405	617	963	1,194	1,860
	(再掲)65-74歳	130	579	386	31	79	160	175	308	504	701	1,121	1,340	2,090
	(再掲)75歳以上	107	616	414	46	74	179	211	303	526	718	1,159	1,609	1,978
	(再掲)20-49歳※	196	465	477	33	54	116	135	210	340	569	938	1,148	2,239
(再掲)40-74歳※	356	480	329	17	52	104	161	269	406	609	934	1,156	1,776	
女性	20-29歳	40	395	246	39	62	104	135	162	347	517	767	785	1,163
	30-39歳	87	450	284	36	98	121	149	237	369	600	866	1,057	1,446
	40-49歳	85	436	393	43	18	43	126	210	345	545	703	1,275	2,879
	50-59歳	97	475	304	29	78	156	188	279	426	568	816	1,085	2,143
	60-69歳	145	569	416	34	8	130	171	286	458	692	1,100	1,341	2,139
	70歳以上	187	526	358	28	21	126	176	274	435	726	1,004	1,152	2,007
	(再掲)20歳以上※	641	486	350	14	40	121	145	252	409	614	895	1,111	2,012
	(再掲)65-74歳	151	573	397	40	18	146	195	294	477	727	1,022	1,323	2,150
	(再掲)75歳以上	137	504	364	37	18	111	169	266	417	608	1,016	1,170	2,136
	(再掲)20-49歳※	212	427	319	22	39	103	136	204	349	547	763	905	1,562
(再掲)40-74歳※	377	510	376	20	39	126	163	261	426	641	949	1,152	2,149	
男女計	20-29歳	82	444	534	49	70	115	133	175	323	517	774	1,132	4,904
	30-39歳	145	463	300	31	96	126	166	235	392	598	904	1,088	1,508
	40-49歳	181	436	346	36	37	64	123	231	355	549	758	1,129	1,617
	50-59歳	181	448	292	26	80	128	174	263	408	532	700	1,008	1,844
	60-69歳	260	549	383	25	48	132	172	292	440	685	1,087	1,293	2,076
	70歳以上	355	559	382	27	37	136	194	293	468	715	1,019	1,348	1,986
	(再掲)20歳以上※	1,204	493	386	14	46	122	152	252	408	616	933	1,150	1,876
	(再掲)65-74歳	281	576	391	26	51	152	193	305	489	715	1,049	1,330	2,151
	(再掲)75歳以上	244	547	387	33	23	126	192	278	459	687	1,070	1,398	1,981
	(再掲)20-49歳※	408	446	407	22	42	113	135	208	348	553	810	1,082	1,564
(再掲)40-74歳※	733	495	354	14	43	121	163	266	412	622	948	1,152	1,958	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

β-カロテン当量(μg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	3,457	4,665	621	93	369	532	1,111	1,922	5,023	8,272	9,611	30,343
	30-39歳	58	3,737	3,091	457	172	315	818	1,248	3,026	4,891	8,904	10,630	12,568
	40-49歳	96	3,620	2,931	420	154	272	532	1,741	2,815	4,891	8,222	10,171	13,030
	50-59歳	84	3,185	2,330	270	101	291	658	1,427	2,897	4,456	5,530	8,010	13,684
	60-69歳	115	4,250	3,277	292	67	209	752	1,754	3,593	6,175	8,353	9,904	17,725
	70歳以上	168	4,996	4,012	317	91	652	984	2,367	4,062	6,751	10,339	12,386	19,477
	(再掲)20歳以上※	563	3,961	3,555	183	100	432	681	1,446	3,005	5,318	8,556	10,206	16,302
	(再掲)65-74歳	130	4,729	3,517	261	84	755	989	2,182	4,001	6,615	9,452	11,314	18,758
	(再掲)75歳以上	107	5,016	4,294	467	84	645	853	2,331	4,028	6,659	10,625	15,356	21,827
	(再掲)20-49歳※	196	3,598	3,633	278	136	328	659	1,241	2,470	4,882	8,474	9,755	16,674
(再掲)40-74歳※	356	3,888	3,035	142	103	291	701	1,757	3,054	5,396	8,221	10,176	13,438	
女性	20-29歳	40	3,139	2,560	461	45	274	413	1,005	2,369	4,874	7,061	8,426	11,303
	30-39歳	87	3,760	3,224	413	147	470	624	1,191	2,767	5,423	8,340	10,425	15,381
	40-49歳	85	3,361	3,534	354	56	225	428	1,005	2,780	4,195	6,306	11,809	20,470
	50-59歳	97	3,827	2,987	232	351	565	1,131	1,888	3,240	4,705	7,202	9,607	19,676
	60-69歳	145	4,858	3,772	300	24	509	1,012	1,994	4,133	6,762	10,346	13,029	16,641
	70歳以上	187	4,392	3,688	302	102	513	732	1,689	3,527	5,915	9,638	11,576	16,120
	(再掲)20歳以上※	641	3,990	3,440	139	95	399	656	1,490	3,088	5,327	8,585	10,563	16,352
	(再掲)65-74歳	151	4,762	3,606	382	73	719	1,055	1,903	4,117	6,754	10,124	10,790	16,481
	(再掲)75歳以上	137	4,224	3,831	409	98	423	652	1,471	3,334	5,838	9,060	12,387	19,966
	(再掲)20-49歳※	212	3,402	3,144	225	54	275	560	1,028	2,594	4,877	7,191	8,882	16,276
(再掲)40-74歳※	377	4,164	3,496	172	80	433	700	1,732	3,253	5,411	8,908	10,563	16,634	
男女計	20-29歳	82	3,304	3,782	382	54	318	606	1,074	2,096	4,872	7,366	9,510	30,343
	30-39歳	145	3,748	3,146	330	162	394	658	1,234	2,913	5,192	8,662	10,354	13,774
	40-49歳	181	3,492	3,236	347	67	273	448	1,389	2,819	4,665	6,622	10,067	16,547
	50-59歳	181	3,506	2,691	211	132	461	737	1,789	2,970	4,486	6,548	8,702	14,469
	60-69歳	260	4,559	3,544	224	68	465	767	1,870	3,721	6,549	9,183	11,025	16,497
	70歳以上	355	4,643	3,833	271	102	535	812	1,967	3,736	6,129	10,123	12,236	17,917
	(再掲)20歳以上※	1,204	3,976	3,494	139	99	406	663	1,469	3,028	5,322	8,584	10,378	16,266
	(再掲)65-74歳	281	4,747	3,560	254	98	747	1,014	2,013	4,035	6,624	9,951	10,919	16,591
	(再掲)75歳以上	244	4,532	4,028	341	97	514	713	1,805	3,556	5,913	10,069	12,522	19,112
	(再掲)20-49歳※	408	3,502	3,401	212	93	328	614	1,206	2,475	4,880	7,495	9,605	15,973
(再掲)40-74歳※	733	4,027	3,276	130	101	423	701	1,738	3,187	5,405	8,416	10,270	16,246	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンD( $\mu\text{g}$ )

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	14.8	18.2	5.06	0.1	0.5	0.9	2.0	4.1	23.3	42.6	55.9	62.2
	30-39歳	58	7.8	9.1	1.13	0.1	0.5	0.9	2.0	3.8	12.0	20.5	28.0	51.4
	40-49歳	96	5.4	6.1	0.53	0.0	0.5	0.6	1.4	3.0	7.9	12.8	18.7	31.1
	50-59歳	84	8.6	10.0	1.24	0.2	0.7	1.2	2.0	4.5	13.0	18.1	32.0	54.2
	60-69歳	115	12.0	12.0	1.17	0.3	0.9	1.4	3.1	8.2	17.2	28.5	34.5	64.7
	70歳以上	168	11.3	9.6	0.65	0.5	1.1	1.8	3.7	8.9	15.8	22.8	30.0	47.4
	(再掲)20歳以上※	563	10.1	11.7	0.96	0.2	0.7	1.1	2.1	5.1	14.2	24.0	33.9	56.2
	(再掲)65-74歳	130	13.4	12.0	1.17	0.3	0.9	1.6	3.8	10.8	19.9	28.8	34.3	67.0
	(再掲)75歳以上	107	11.4	9.8	0.79	0.4	1.4	2.3	3.7	9.0	16.0	24.3	32.9	47.3
	(再掲)20-49歳※	196	9.3	12.8	1.82	0.0	0.6	0.8	1.8	3.5	12.4	24.5	33.8	59.7
(再掲)40-74歳※	356	9.0	10.0	0.59	0.3	0.6	1.1	1.9	4.8	13.4	22.0	30.7	44.4	
女性	20-29歳	40	10.6	15.0	4.74	0.1	0.2	0.7	1.8	3.7	11.6	37.6	48.2	51.9
	30-39歳	87	7.2	8.7	0.91	0.2	0.4	0.8	1.4	3.1	11.6	22.2	26.0	36.6
	40-49歳	85	5.9	6.6	0.66	0.4	0.5	1.1	1.7	2.9	8.3	13.6	17.0	45.2
	50-59歳	97	8.6	10.5	1.18	0.2	0.5	0.9	1.9	4.0	11.7	27.7	33.5	49.7
	60-69歳	145	11.2	10.8	1.01	0.3	0.8	1.3	2.6	8.8	15.4	24.5	31.2	56.4
	70歳以上	187	9.6	8.1	0.56	0.3	0.9	1.5	3.1	7.6	14.1	19.8	26.0	42.2
	(再掲)20歳以上※	641	9.0	10.1	0.79	0.2	0.7	1.0	2.1	5.2	13.0	21.1	30.7	50.5
	(再掲)65-74歳	151	11.5	10.4	0.73	0.6	0.9	1.3	2.9	9.1	15.7	26.9	30.4	55.0
	(再掲)75歳以上	137	9.8	8.1	0.60	0.1	1.0	1.7	3.7	7.8	14.3	19.9	25.3	46.7
	(再掲)20-49歳※	212	7.8	10.7	1.60	0.1	0.5	0.9	1.5	3.1	10.3	19.0	33.3	50.2
(再掲)40-74歳※	377	8.7	9.5	0.51	0.3	0.7	1.0	2.0	5.0	12.7	20.7	30.4	45.5	
男女計	20-29歳	82	12.8	16.8	4.78	0.1	0.3	0.9	2.0	4.0	14.6	43.5	52.7	62.2
	30-39歳	145	7.5	8.9	0.85	0.2	0.5	0.8	1.5	3.3	11.9	20.4	26.6	42.9
	40-49歳	181	5.6	6.3	0.50	0.0	0.5	0.7	1.5	3.0	8.2	13.5	16.9	31.6
	50-59歳	181	8.6	10.2	1.17	0.2	0.6	1.0	2.0	4.3	12.5	20.9	33.9	49.6
	60-69歳	260	11.6	11.4	0.98	0.3	0.9	1.3	2.8	8.4	16.1	27.2	34.2	58.8
	70歳以上	355	10.3	8.8	0.51	0.4	0.9	1.6	3.1	8.0	14.7	21.0	27.6	45.8
	(再掲)20歳以上※	1,204	9.5	10.9	0.85	0.2	0.7	1.0	2.1	5.1	13.5	23.0	33.4	52.5
	(再掲)65-74歳	281	12.4	11.2	0.85	0.5	0.9	1.5	3.3	9.7	17.0	27.4	31.1	58.9
	(再掲)75歳以上	244	10.4	8.8	0.57	0.4	1.1	2.0	3.7	8.0	14.7	20.9	27.2	46.1
	(再掲)20-49歳※	408	8.6	11.9	1.69	0.1	0.5	0.8	1.7	3.3	11.6	23.0	33.4	53.3
(再掲)40-74歳※	733	8.9	9.7	0.52	0.3	0.7	1.0	1.9	4.9	13.1	21.1	30.4	44.9	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンE(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	6.1	2.8	0.33	2.0	2.3	2.6	4.1	6.1	7.6	10.1	11.0	15.8
	30-39歳	58	6.7	2.8	0.42	0.8	2.2	2.8	4.5	6.6	8.5	10.4	12.0	12.8
	40-49歳	96	6.3	2.6	0.27	0.8	2.5	3.2	4.4	6.1	8.1	9.5	11.4	13.6
	50-59歳	84	6.7	3.6	0.39	1.3	2.7	3.8	4.3	5.6	8.1	10.8	15.4	23.1
	60-69歳	115	7.5	3.4	0.32	1.2	2.7	3.4	4.6	7.4	9.5	12.5	14.2	17.1
	70歳以上	168	7.6	3.7	0.36	1.9	2.7	3.7	4.7	7.2	9.5	11.8	13.0	23.5
	(再掲)20歳以上※	563	6.9	3.2	0.18	1.5	2.6	3.2	4.5	6.5	8.6	11.0	12.5	17.2
	(再掲)65-74歳	130	7.5	3.3	0.31	2.2	2.7	3.8	4.9	7.2	9.5	11.6	14.0	17.8
	(再掲)75歳以上	107	7.6	3.8	0.44	1.3	2.7	3.6	4.7	7.1	9.4	11.9	12.8	26.4
	(再掲)20-49歳※	196	6.4	2.7	0.19	1.5	2.5	2.7	4.2	6.1	8.2	9.9	11.2	14.1
(再掲)40-74歳※	356	6.9	3.3	0.17	1.4	2.7	3.6	4.5	6.4	8.8	11.1	13.4	17.3	
女性	20-29歳	40	5.8	2.4	0.50	1.1	1.9	2.5	4.4	5.7	7.0	9.2	11.4	11.7
	30-39歳	87	6.5	3.0	0.42	1.4	2.4	2.8	4.2	6.1	7.9	9.9	12.4	17.2
	40-49歳	85	6.0	2.9	0.20	1.7	2.3	2.6	3.9	5.5	7.4	9.1	12.0	15.8
	50-59歳	97	7.1	3.1	0.30	1.8	3.3	4.0	4.6	6.7	8.8	10.2	13.9	17.1
	60-69歳	145	6.9	3.7	0.26	1.2	2.6	3.1	4.2	6.0	8.8	12.4	14.5	20.8
	70歳以上	187	6.5	3.4	0.31	1.1	2.0	2.8	4.1	6.2	8.2	10.5	12.6	19.4
	(再掲)20歳以上※	641	6.5	3.2	0.17	1.4	2.4	2.9	4.2	6.0	8.1	10.1	12.4	17.1
	(再掲)65-74歳	151	7.0	3.7	0.32	1.3	2.4	2.9	4.2	6.3	8.8	11.5	14.0	20.6
	(再掲)75歳以上	137	6.2	3.3	0.36	1.1	1.8	2.7	3.9	5.8	7.9	9.7	12.5	19.1
	(再掲)20-49歳※	212	6.1	2.8	0.20	1.3	2.4	2.7	4.1	5.8	7.5	9.3	11.5	15.7
(再掲)40-74歳※	377	6.7	3.3	0.18	1.5	2.4	3.2	4.3	6.2	8.5	10.6	13.6	17.3	
男女計	20-29歳	82	6.0	2.6	0.28	1.2	2.3	2.6	4.2	5.9	7.0	9.3	11.3	15.8
	30-39歳	145	6.6	2.9	0.36	1.1	2.4	2.8	4.4	6.3	8.3	10.1	12.2	15.3
	40-49歳	181	6.1	2.8	0.18	1.4	2.4	2.9	4.1	5.7	7.7	9.3	11.0	15.6
	50-59歳	181	6.9	3.3	0.25	1.8	3.1	3.9	4.5	6.3	8.6	10.4	14.4	17.3
	60-69歳	260	7.2	3.5	0.21	1.3	2.7	3.2	4.4	6.6	9.1	12.4	14.3	17.7
	70歳以上	355	6.9	3.6	0.30	1.3	2.3	3.0	4.4	6.5	8.7	11.1	12.8	19.6
	(再掲)20歳以上※	1,204	6.7	3.2	0.16	1.4	2.4	3.0	4.4	6.2	8.4	10.6	12.5	17.2
	(再掲)65-74歳	281	7.2	3.5	0.28	1.4	2.6	3.2	4.6	6.6	9.1	11.5	14.1	18.9
	(再掲)75歳以上	244	6.7	3.6	0.36	1.1	2.2	2.9	4.4	6.2	8.6	11.1	12.6	22.1
	(再掲)20-49歳※	408	6.2	2.7	0.17	1.4	2.4	2.7	4.2	6.0	7.8	9.7	11.3	15.7
(再掲)40-74歳※	733	6.8	3.3	0.15	1.4	2.6	3.3	4.4	6.3	8.6	10.8	13.5	17.2	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンK( $\mu\text{g}$ )

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	211	130	23.3	36	51	70	114	185	293	375	513	569
	30-39歳	58	335	241	39.2	18	41	53	132	304	469	714	854	1,056
	40-49歳	96	223	138	14.6	4	48	80	122	188	316	402	480	872
	50-59歳	84	295	179	19.6	25	55	72	162	283	393	532	616	888
	60-69歳	115	304	190	20.1	17	61	81	148	265	452	523	674	874
	70歳以上	168	313	214	22.5	32	53	73	159	277	425	641	693	1,165
	(再掲)20歳以上※	563	280	191	14.3	28	53	75	129	239	374	516	645	864
	(再掲)65-74歳	130	315	193	21.8	17	68	88	162	263	455	640	693	769
	(再掲)75歳以上	107	305	222	29.6	32	43	64	146	277	420	547	658	1,280
	(再掲)20-49歳※	196	251	179	17.5	18	48	71	122	204	342	492	599	868
(再掲)40-74歳※	356	280	178	13.6	21	59	79	143	239	392	516	641	791	
女性	20-29歳	40	196	170	29.9	8	35	58	86	144	237	397	536	964
	30-39歳	87	241	170	21.5	24	50	67	111	201	326	483	584	991
	40-49歳	85	217	124	8.8	43	69	85	124	183	281	392	451	664
	50-59歳	97	253	149	19.6	21	57	83	125	235	351	472	550	674
	60-69歳	145	311	207	24.1	5	67	84	146	257	426	627	690	912
	70歳以上	187	254	192	15.7	25	49	64	112	238	326	472	628	1,133
	(再掲)20歳以上※	641	248	178	9.8	23	54	75	114	210	334	474	594	886
	(再掲)65-74歳	151	276	193	20.7	9	54	83	128	224	363	598	650	896
	(再掲)75歳以上	137	262	205	17.9	21	45	62	113	238	338	476	627	1,299
	(再掲)20-49歳※	212	217	155	13.4	14	56	71	110	185	282	407	513	933
(再掲)40-74歳※	377	258	167	11.6	30	67	83	126	214	349	479	621	803	
男女計	20-29歳	82	204	150	16.6	10	48	69	97	168	265	377	512	964
	30-39歳	145	289	214	27.1	22	45	65	122	233	385	593	737	1,024
	40-49歳	181	220	131	9.5	20	63	84	124	184	292	398	472	669
	50-59歳	181	274	165	16.6	26	57	79	136	255	371	498	565	814
	60-69歳	260	307	198	18.0	16	62	85	148	263	439	597	670	862
	70歳以上	355	279	203	17.4	32	50	70	124	242	364	513	650	1,133
	(再掲)20歳以上※	1,204	264	185	10.7	24	54	76	124	221	354	499	625	874
	(再掲)65-74歳	281	294	193	17.7	21	67	87	134	241	412	613	666	817
	(再掲)75歳以上	244	279	213	19.6	27	45	64	118	241	367	489	643	1,260
	(再掲)20-49歳※	408	234	168	11.3	17	51	71	114	194	318	452	563	881
(再掲)40-74歳※	733	269	173	11.4	27	61	82	130	224	370	504	626	792	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンB<sub>1</sub>(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	0.78	0.33	0.034	0.18	0.30	0.40	0.64	0.74	0.86	1.33	1.53	1.92
	30-39歳	58	1.02	0.48	0.052	0.22	0.36	0.51	0.68	0.97	1.27	1.45	1.63	3.57
	40-49歳	96	0.97	0.39	0.041	0.29	0.43	0.51	0.68	0.93	1.14	1.49	1.77	2.31
	50-59歳	84	0.94	0.42	0.060	0.28	0.47	0.54	0.70	0.82	1.08	1.57	1.81	2.66
	60-69歳	115	0.99	0.40	0.032	0.34	0.47	0.58	0.71	0.94	1.16	1.41	1.81	2.67
	70歳以上	168	0.92	0.36	0.029	0.31	0.42	0.48	0.66	0.87	1.10	1.38	1.72	2.03
	(再掲)20歳以上※	563	0.94	0.40	0.022	0.25	0.42	0.51	0.66	0.86	1.13	1.40	1.70	2.30
	(再掲)65-74歳	130	1.01	0.39	0.038	0.41	0.49	0.57	0.73	0.99	1.14	1.44	1.82	2.53
	(再掲)75歳以上	107	0.89	0.34	0.044	0.26	0.40	0.47	0.66	0.81	1.08	1.34	1.50	1.94
	(再掲)20-49歳※	196	0.92	0.41	0.029	0.22	0.39	0.45	0.65	0.86	1.16	1.40	1.59	2.31
(再掲)40-74歳※	356	0.97	0.40	0.025	0.34	0.45	0.53	0.70	0.89	1.13	1.46	1.79	2.43	
女性	20-29歳	40	0.66	0.25	0.040	0.20	0.28	0.36	0.52	0.62	0.76	1.03	1.34	1.38
	30-39歳	87	0.79	0.31	0.041	0.28	0.38	0.43	0.56	0.76	1.03	1.23	1.45	1.71
	40-49歳	85	0.80	0.33	0.037	0.30	0.39	0.42	0.53	0.76	1.04	1.27	1.29	2.16
	50-59歳	97	0.77	0.27	0.024	0.24	0.41	0.46	0.56	0.73	0.90	1.19	1.28	1.82
	60-69歳	145	0.85	0.34	0.029	0.14	0.38	0.49	0.61	0.82	1.06	1.23	1.38	1.97
	70歳以上	187	0.76	0.31	0.029	0.19	0.35	0.44	0.56	0.71	0.87	1.20	1.35	1.88
	(再掲)20歳以上※	641	0.78	0.31	0.019	0.23	0.36	0.44	0.56	0.71	0.93	1.20	1.30	1.85
	(再掲)65-74歳	151	0.84	0.33	0.031	0.16	0.36	0.47	0.60	0.85	1.04	1.24	1.37	1.96
	(再掲)75歳以上	137	0.74	0.32	0.034	0.18	0.32	0.44	0.54	0.69	0.87	1.12	1.42	1.99
	(再掲)20-49歳※	212	0.75	0.31	0.029	0.25	0.35	0.41	0.53	0.69	0.91	1.23	1.32	1.73
(再掲)40-74歳※	377	0.81	0.31	0.019	0.24	0.39	0.46	0.57	0.78	1.00	1.23	1.29	1.90	
男女計	20-29歳	82	0.72	0.30	0.032	0.18	0.30	0.38	0.56	0.66	0.82	1.11	1.37	1.92
	30-39歳	145	0.91	0.42	0.037	0.23	0.38	0.45	0.62	0.85	1.18	1.37	1.55	2.86
	40-49歳	181	0.88	0.37	0.033	0.31	0.39	0.47	0.58	0.85	1.11	1.29	1.70	2.20
	50-59歳	181	0.86	0.36	0.030	0.24	0.43	0.51	0.64	0.79	0.95	1.29	1.57	2.62
	60-69歳	260	0.92	0.38	0.020	0.21	0.43	0.54	0.65	0.89	1.11	1.29	1.73	2.25
	70歳以上	355	0.83	0.34	0.026	0.24	0.38	0.45	0.59	0.77	1.00	1.29	1.49	1.98
	(再掲)20歳以上※	1,204	0.85	0.36	0.018	0.24	0.39	0.46	0.62	0.79	1.04	1.30	1.53	1.99
	(再掲)65-74歳	281	0.92	0.37	0.025	0.23	0.43	0.51	0.66	0.89	1.09	1.29	1.67	2.10
	(再掲)75歳以上	244	0.80	0.33	0.032	0.20	0.35	0.44	0.58	0.74	0.93	1.25	1.45	1.96
	(再掲)20-49歳※	408	0.84	0.37	0.026	0.23	0.35	0.43	0.58	0.77	1.04	1.31	1.51	1.87
(再掲)40-74歳※	733	0.89	0.37	0.018	0.29	0.43	0.49	0.63	0.85	1.09	1.29	1.59	2.12	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンB<sub>2</sub>(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	1.07	0.41	0.078	0.38	0.42	0.47	0.76	1.03	1.36	1.66	1.73	1.93
	30-39歳	58	1.30	0.59	0.051	0.19	0.53	0.61	1.00	1.25	1.56	1.92	2.31	4.48
	40-49歳	96	1.01	0.42	0.040	0.20	0.37	0.45	0.71	0.99	1.19	1.55	1.93	2.15
	50-59歳	84	1.22	0.48	0.055	0.38	0.56	0.64	0.88	1.16	1.54	1.81	2.26	3.03
	60-69歳	115	1.31	0.46	0.040	0.42	0.62	0.76	0.99	1.24	1.58	1.91	2.23	2.60
	70歳以上	168	1.33	0.47	0.046	0.38	0.60	0.75	1.00	1.30	1.58	2.10	2.21	2.43
	(再掲)20歳以上※	563	1.21	0.49	0.024	0.35	0.51	0.63	0.89	1.17	1.49	1.83	2.11	2.49
	(再掲)65-74歳	130	1.38	0.46	0.040	0.42	0.62	0.81	1.05	1.36	1.63	2.09	2.24	2.59
	(再掲)75歳以上	107	1.32	0.48	0.058	0.33	0.60	0.72	0.98	1.30	1.55	2.11	2.27	2.51
	(再掲)20-49歳※	196	1.11	0.48	0.038	0.24	0.42	0.53	0.77	1.08	1.39	1.67	1.92	2.52
(再掲)40-74歳※	356	1.20	0.47	0.026	0.32	0.55	0.64	0.86	1.16	1.50	1.85	2.10	2.48	
女性	20-29歳	40	1.00	0.39	0.071	0.27	0.30	0.47	0.67	1.02	1.28	1.43	1.75	2.05
	30-39歳	87	0.98	0.36	0.049	0.26	0.46	0.52	0.70	0.96	1.17	1.50	1.55	2.15
	40-49歳	85	1.03	0.47	0.061	0.37	0.44	0.54	0.72	0.98	1.16	1.61	1.94	3.17
	50-59歳	97	1.13	0.40	0.051	0.25	0.54	0.58	0.88	1.13	1.27	1.61	1.91	2.47
	60-69歳	145	1.25	0.59	0.056	0.31	0.53	0.64	0.91	1.16	1.51	1.81	2.11	3.83
	70歳以上	187	1.16	0.43	0.040	0.30	0.54	0.64	0.87	1.15	1.39	1.70	1.93	2.42
	(再掲)20歳以上※	641	1.11	0.46	0.021	0.30	0.49	0.57	0.82	1.06	1.34	1.65	1.88	2.68
	(再掲)65-74歳	151	1.24	0.55	0.048	0.32	0.54	0.68	0.89	1.20	1.50	1.79	2.00	3.65
	(再掲)75歳以上	137	1.17	0.44	0.044	0.30	0.53	0.59	0.89	1.14	1.37	1.73	1.99	2.60
	(再掲)20-49歳※	212	1.01	0.41	0.032	0.27	0.44	0.52	0.70	1.00	1.20	1.49	1.74	2.70
(再掲)40-74歳※	377	1.14	0.49	0.025	0.34	0.53	0.60	0.84	1.09	1.34	1.71	1.93	2.76	
男女計	20-29歳	82	1.04	0.40	0.059	0.27	0.41	0.48	0.73	1.02	1.34	1.62	1.72	2.05
	30-39歳	145	1.14	0.51	0.040	0.23	0.50	0.56	0.81	1.09	1.41	1.65	1.98	3.44
	40-49歳	181	1.02	0.45	0.039	0.25	0.40	0.53	0.71	0.98	1.18	1.55	1.92	2.74
	50-59歳	181	1.17	0.44	0.039	0.28	0.55	0.59	0.89	1.15	1.42	1.73	2.01	2.48
	60-69歳	260	1.28	0.53	0.035	0.40	0.58	0.72	0.94	1.21	1.54	1.89	2.18	3.12
	70歳以上	355	1.23	0.45	0.035	0.33	0.55	0.68	0.91	1.22	1.48	1.88	2.14	2.37
	(再掲)20歳以上※	1,204	1.16	0.47	0.019	0.31	0.49	0.60	0.84	1.11	1.40	1.73	2.02	2.52
	(再掲)65-74歳	281	1.30	0.52	0.038	0.39	0.59	0.73	0.98	1.24	1.58	1.93	2.14	2.91
	(再掲)75歳以上	244	1.23	0.47	0.039	0.31	0.54	0.63	0.90	1.22	1.44	1.88	2.16	2.51
	(再掲)20-49歳※	408	1.06	0.45	0.027	0.27	0.43	0.53	0.74	1.03	1.30	1.61	1.79	2.58
(再掲)40-74歳※	733	1.17	0.48	0.021	0.35	0.53	0.61	0.85	1.11	1.42	1.77	2.05	2.66	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ナイアシン(mgNE)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	13.8	6.4	0.98	3.1	5.3	6.7	8.3	12.7	17.4	24.7	25.2	25.8
	30-39歳	58	17.6	7.4	1.05	4.5	6.9	9.3	11.6	16.5	21.9	27.8	30.5	44.4
	40-49歳	96	15.0	6.0	0.64	4.6	6.4	7.6	9.8	14.8	18.5	23.6	26.4	33.3
	50-59歳	84	17.0	7.3	0.82	6.7	7.1	9.0	11.5	14.8	22.7	28.1	31.3	38.5
	60-69歳	115	18.5	9.6	1.08	5.3	8.8	9.4	12.3	16.5	21.1	29.2	37.2	55.7
	70歳以上	168	16.4	6.9	0.61	5.1	7.7	8.4	11.5	15.0	20.3	25.1	28.5	41.1
	(再掲)20歳以上※	563	16.4	7.5	0.42	5.1	6.9	8.0	11.1	14.9	20.6	25.3	29.2	44.5
	(再掲)65-74歳	130	18.8	9.2	1.18	5.3	8.6	9.8	13.4	17.4	21.6	28.2	32.5	55.7
	(再掲)75歳以上	107	15.7	7.0	0.82	4.9	6.6	8.1	10.8	14.4	18.5	24.5	26.9	44.2
	(再掲)20-49歳※	196	15.3	6.7	0.53	3.6	6.3	7.6	10.4	14.5	19.4	24.7	26.2	36.3
(再掲)40-74歳※	356	16.9	7.7	0.42	5.4	7.4	8.6	11.7	15.6	20.8	26.4	30.9	49.6	
女性	20-29歳	40	14.1	7.0	1.63	2.5	4.9	5.4	8.4	12.5	19.3	23.3	25.6	32.6
	30-39歳	87	12.5	4.8	0.57	4.8	5.5	7.0	9.2	11.5	15.6	19.7	20.9	27.6
	40-49歳	85	12.7	4.7	0.39	5.0	5.5	6.8	8.8	12.2	15.2	18.6	22.0	26.9
	50-59歳	97	14.1	5.9	0.65	3.8	6.2	7.0	10.0	12.5	18.2	23.5	24.7	30.4
	60-69歳	145	14.9	5.9	0.51	2.7	6.3	7.9	9.9	14.2	18.9	22.6	24.5	32.0
	70歳以上	187	13.5	5.3	0.51	4.7	6.2	7.1	9.3	13.1	16.3	21.2	22.9	33.0
	(再掲)20歳以上※	641	13.6	5.7	0.40	4.4	5.7	6.9	9.5	12.9	17.2	22.0	23.8	30.3
	(再掲)65-74歳	151	14.9	5.6	0.40	2.7	6.4	8.0	10.7	14.2	18.8	22.5	23.8	32.3
	(再掲)75歳以上	137	13.2	5.3	0.57	4.3	6.1	7.1	8.9	13.0	16.0	20.3	23.4	33.8
	(再掲)20-49歳※	212	13.1	5.6	0.62	4.2	5.4	6.5	9.1	11.8	16.7	21.1	23.2	27.8
(再掲)40-74歳※	377	13.9	5.5	0.29	4.5	6.2	7.4	10.0	13.2	17.4	22.1	24.0	30.2	
男女計	20-29歳	82	13.9	6.7	1.02	2.7	5.2	6.4	8.4	12.6	18.3	24.6	25.8	32.6
	30-39歳	145	15.1	6.8	0.66	4.8	5.8	7.7	10.3	14.4	19.1	24.3	28.3	41.2
	40-49歳	181	13.8	5.5	0.47	5.0	6.0	7.2	9.7	13.2	17.0	21.0	24.8	30.1
	50-59歳	181	15.6	6.8	0.60	5.1	6.7	7.9	10.6	13.6	19.8	24.6	30.2	35.6
	60-69歳	260	16.7	8.1	0.71	4.4	7.6	8.8	11.5	15.6	20.1	24.8	30.9	54.0
	70歳以上	355	14.7	6.2	0.49	5.0	6.3	7.7	10.2	13.8	18.1	22.9	25.5	33.9
	(再掲)20歳以上※	1,204	14.9	6.7	0.34	4.5	6.3	7.5	10.3	13.8	18.4	23.7	26.0	33.9
	(再掲)65-74歳	281	16.7	7.8	0.68	4.5	7.5	9.0	11.7	15.7	20.4	24.2	28.4	53.1
	(再掲)75歳以上	244	14.2	6.1	0.59	4.7	6.2	7.5	9.5	13.7	17.8	22.7	25.0	35.0
	(再掲)20-49歳※	408	14.2	6.3	0.43	4.1	5.6	6.9	9.7	13.0	17.9	23.7	25.3	32.3
(再掲)40-74歳※	733	15.4	6.9	0.30	5.0	6.7	7.8	10.7	14.2	18.9	23.8	27.1	35.9	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンB<sub>6</sub>(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	1.10	0.49	0.057	0.31	0.45	0.58	0.75	0.97	1.55	1.63	2.14	2.43
	30-39歳	58	1.32	0.57	0.089	0.26	0.44	0.60	0.99	1.33	1.51	2.04	2.55	3.14
	40-49歳	96	1.14	0.41	0.045	0.15	0.52	0.59	0.82	1.16	1.44	1.59	1.80	2.44
	50-59歳	84	1.24	0.55	0.053	0.28	0.62	0.67	0.88	1.06	1.48	2.04	2.43	3.87
	60-69歳	115	1.36	0.59	0.058	0.26	0.71	0.75	1.00	1.28	1.58	2.03	2.60	3.68
	70歳以上	168	1.36	0.51	0.052	0.45	0.60	0.74	1.00	1.32	1.69	2.04	2.36	2.87
	(再掲)20歳以上※	563	1.26	0.53	0.033	0.31	0.57	0.67	0.87	1.22	1.54	1.86	2.33	2.88
	(再掲)65-74歳	130	1.44	0.56	0.065	0.36	0.71	0.84	1.08	1.37	1.68	2.06	2.71	3.58
	(再掲)75歳以上	107	1.34	0.53	0.063	0.45	0.55	0.67	0.97	1.28	1.69	2.07	2.36	3.08
	(再掲)20-49歳※	196	1.18	0.49	0.039	0.28	0.52	0.59	0.79	1.16	1.49	1.77	2.17	2.72
(再掲)40-74歳※	356	1.27	0.52	0.032	0.29	0.60	0.72	0.91	1.21	1.52	1.86	2.37	2.88	
女性	20-29歳	40	0.93	0.38	0.061	0.17	0.40	0.51	0.69	0.87	1.17	1.37	1.61	2.41
	30-39歳	87	0.99	0.35	0.041	0.36	0.46	0.57	0.76	0.95	1.18	1.47	1.73	1.99
	40-49歳	85	1.01	0.41	0.045	0.37	0.44	0.56	0.69	0.97	1.27	1.55	1.79	2.18
	50-59歳	97	1.07	0.38	0.043	0.22	0.50	0.63	0.80	1.04	1.23	1.58	1.81	2.37
	60-69歳	145	1.23	0.47	0.042	0.32	0.54	0.68	0.91	1.12	1.50	1.84	2.32	2.61
	70歳以上	187	1.10	0.41	0.028	0.35	0.49	0.62	0.82	1.08	1.32	1.57	1.85	2.23
	(再掲)20歳以上※	641	1.07	0.42	0.023	0.32	0.47	0.59	0.77	1.02	1.28	1.59	1.85	2.33
	(再掲)65-74歳	151	1.22	0.44	0.038	0.32	0.59	0.70	0.93	1.17	1.43	1.80	2.25	2.48
	(再掲)75歳以上	137	1.09	0.42	0.035	0.34	0.45	0.58	0.81	1.08	1.32	1.58	1.85	2.59
	(再掲)20-49歳※	212	0.98	0.38	0.034	0.23	0.43	0.55	0.72	0.90	1.19	1.45	1.68	2.19
(再掲)40-74歳※	377	1.11	0.43	0.024	0.35	0.53	0.62	0.81	1.06	1.33	1.67	1.95	2.35	
男女計	20-29歳	82	1.02	0.45	0.045	0.19	0.42	0.57	0.72	0.89	1.26	1.60	1.85	2.43
	30-39歳	145	1.16	0.50	0.054	0.29	0.45	0.59	0.78	1.10	1.42	1.78	2.03	2.95
	40-49歳	181	1.08	0.41	0.039	0.33	0.52	0.58	0.76	1.08	1.34	1.58	1.83	2.35
	50-59歳	181	1.16	0.48	0.035	0.27	0.58	0.64	0.88	1.06	1.35	1.75	2.21	2.60
	60-69歳	260	1.29	0.54	0.044	0.32	0.61	0.73	0.94	1.23	1.54	1.95	2.33	3.02
	70歳以上	355	1.21	0.47	0.034	0.41	0.55	0.66	0.90	1.17	1.44	1.84	2.07	2.76
	(再掲)20歳以上※	1,204	1.16	0.48	0.026	0.32	0.52	0.63	0.82	1.09	1.42	1.78	2.04	2.71
	(再掲)65-74歳	281	1.32	0.52	0.044	0.35	0.65	0.74	0.97	1.26	1.54	1.95	2.32	2.95
	(再掲)75歳以上	244	1.19	0.48	0.040	0.36	0.50	0.62	0.84	1.12	1.39	1.84	2.07	2.84
	(再掲)20-49歳※	408	1.08	0.45	0.031	0.26	0.46	0.58	0.76	1.01	1.35	1.62	1.90	2.46
(再掲)40-74歳※	733	1.19	0.48	0.025	0.33	0.57	0.64	0.87	1.13	1.44	1.79	2.09	2.73	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンB<sub>12</sub>( $\mu$ g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	6.00	6.74	1.02	0.47	0.64	1.22	2.47	3.98	8.05	13.06	20.85	40.12
	30-39歳	58	6.24	6.31	0.72	0.22	0.83	0.90	2.40	4.69	7.59	14.01	22.95	35.22
	40-49歳	96	4.62	4.55	0.51	0.09	0.85	1.16	1.76	3.35	5.51	10.89	13.98	25.21
	50-59歳	84	6.69	5.91	1.07	0.46	0.71	1.19	2.55	4.94	8.72	16.27	22.67	24.75
	60-69歳	115	9.17	11.33	1.19	0.40	1.03	1.23	2.96	5.67	10.53	23.90	27.70	71.03
	70歳以上	168	7.71	5.90	0.41	0.31	1.30	1.95	3.58	6.23	9.93	14.97	18.78	29.82
	(再掲)20歳以上※	563	6.83	7.27	0.37	0.36	0.88	1.31	2.57	4.52	8.81	14.75	21.71	35.99
	(再掲)65-74歳	130	9.51	10.91	1.18	0.73	1.04	1.74	3.48	6.57	10.71	20.20	27.64	66.79
	(再掲)75歳以上	107	8.04	6.45	0.56	0.15	1.38	1.93	3.58	6.32	10.38	16.19	25.14	33.28
	(再掲)20-49歳※	196	5.55	5.89	0.42	0.28	0.81	1.12	2.02	4.00	7.07	11.24	16.43	35.96
(再掲)40-74歳※	356	6.88	7.78	0.55	0.38	0.99	1.23	2.38	4.57	8.81	15.73	21.85	36.89	
女性	20-29歳	40	6.26	7.11	1.69	0.13	0.73	1.01	1.42	3.56	8.76	20.08	24.96	26.24
	30-39歳	87	5.13	6.90	0.69	0.17	0.62	0.78	1.46	2.31	5.67	13.80	20.03	35.75
	40-49歳	85	4.29	4.02	0.43	0.40	0.80	1.01	2.01	2.76	5.23	8.93	13.61	24.51
	50-59歳	97	6.09	6.79	0.97	0.41	0.54	1.27	1.83	3.92	6.65	16.84	24.33	35.80
	60-69歳	145	7.39	8.30	0.78	0.55	1.04	1.28	2.60	4.95	8.92	16.24	21.84	50.70
	70歳以上	187	6.62	5.54	0.41	0.22	1.01	1.23	2.78	5.35	8.91	13.23	17.38	28.38
	(再掲)20歳以上※	641	6.09	6.50	0.50	0.42	0.78	1.10	2.00	4.13	8.01	13.53	19.31	29.42
	(再掲)65-74歳	151	7.75	8.19	0.68	0.46	1.05	1.24	2.49	5.36	9.77	17.62	23.15	47.02
	(再掲)75歳以上	137	6.50	5.27	0.41	0.46	1.04	1.39	3.10	5.36	8.76	13.22	15.69	32.31
	(再掲)20-49歳※	212	5.18	6.07	0.79	0.39	0.74	1.00	1.61	2.79	6.36	12.15	19.22	29.88
(再掲)40-74歳※	377	6.10	6.69	0.46	0.45	0.88	1.20	2.05	4.14	7.67	13.84	20.12	28.16	
男女計	20-29歳	82	6.12	6.88	0.99	0.19	0.71	1.03	2.01	3.87	8.41	13.29	24.78	40.12
	30-39歳	145	5.70	6.61	0.61	0.23	0.75	0.85	1.71	3.43	7.21	13.81	19.92	35.26
	40-49歳	181	4.46	4.29	0.40	0.30	0.86	1.11	1.85	2.98	5.25	10.47	13.84	24.79
	50-59歳	181	6.39	6.35	0.96	0.45	0.70	1.26	2.20	4.47	7.54	16.43	22.42	27.39
	60-69歳	260	8.26	9.93	0.91	0.52	1.05	1.25	2.88	5.19	9.83	18.56	25.25	58.92
	70歳以上	355	7.07	5.71	0.35	0.31	1.05	1.49	3.21	5.64	9.47	14.47	17.81	27.27
	(再掲)20歳以上※	1,204	6.45	6.89	0.40	0.39	0.85	1.20	2.16	4.28	8.39	13.89	19.90	33.86
	(再掲)65-74歳	281	8.57	9.58	0.82	0.56	1.05	1.35	2.96	5.67	10.05	18.64	24.84	52.32
	(再掲)75歳以上	244	7.10	5.80	0.40	0.31	1.07	1.55	3.45	5.70	9.25	13.63	17.61	31.52
	(再掲)20-49歳※	408	5.37	5.98	0.47	0.33	0.76	1.02	1.86	3.47	6.63	11.57	17.66	30.42
(再掲)40-74歳※	733	6.49	7.26	0.47	0.44	0.90	1.24	2.15	4.28	8.39	14.86	20.55	33.30	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

葉酸(μg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	235	109	21.1	65	69	114	167	201	329	373	480	487
	30-39歳	58	318	151	16.7	43	111	143	222	297	397	504	519	1,008
	40-49歳	96	258	98	10.9	39	103	140	198	232	328	381	438	493
	50-59歳	84	309	148	16.7	85	138	162	204	259	386	516	598	1,006
	60-69歳	115	382	183	19.2	132	169	177	246	358	464	602	733	1,131
	70歳以上	168	370	150	14.2	155	168	197	260	349	440	562	656	895
	(再掲)20歳以上※	563	316	153	10.8	67	125	160	206	291	394	503	580	852
	(再掲)65-74歳	130	400	183	21.5	154	172	188	264	375	470	635	728	1,121
	(再掲)75歳以上	107	360	151	16.9	155	167	189	260	340	426	558	659	944
	(再掲)20-49歳※	196	267	123	11.2	55	99	125	178	240	339	436	492	597
(再掲)40-74歳※	356	327	157	9.0	96	143	170	211	303	406	516	610	958	
女性	20-29歳	40	242	125	26.7	61	78	111	137	232	322	377	380	818
	30-39歳	87	255	112	12.7	81	103	131	171	237	317	398	489	704
	40-49歳	85	253	93	16.3	91	119	137	180	239	311	390	418	486
	50-59歳	97	314	116	13.8	75	141	161	230	317	373	458	524	671
	60-69歳	145	352	149	10.2	72	134	184	236	328	440	546	644	758
	70歳以上	187	334	144	9.5	93	154	184	247	314	394	506	621	888
	(再掲)20歳以上※	641	300	135	8.0	77	119	142	209	286	373	460	541	725
	(再掲)65-74歳	151	356	141	13.9	72	161	196	261	335	439	524	622	794
	(再掲)75歳以上	137	329	150	12.1	89	147	171	245	301	387	496	628	1,043
	(再掲)20-49歳※	212	250	109	11.7	64	102	122	169	236	313	378	416	709
(再掲)40-74歳※	377	313	129	7.8	90	136	157	221	299	389	480	544	705	
男女計	20-29歳	82	238	116	19.6	61	69	118	153	224	323	376	444	818
	30-39歳	145	287	136	11.3	50	113	132	186	266	370	492	505	833
	40-49歳	181	256	95	12.4	84	108	138	183	236	320	387	434	490
	50-59歳	181	311	133	12.3	81	141	164	215	285	382	507	582	700
	60-69歳	260	367	167	10.5	84	158	178	245	347	452	551	672	1,027
	70歳以上	355	349	147	10.7	100	163	191	247	330	411	530	633	849
	(再掲)20歳以上※	1,204	308	144	8.4	69	120	155	208	289	381	487	561	800
	(再掲)65-74歳	281	377	164	15.8	111	169	194	262	362	452	564	694	1,000
	(再掲)75歳以上	244	341	151	11.7	96	158	182	247	319	405	529	629	956
	(再掲)20-49歳※	408	259	117	9.7	63	101	125	173	238	327	397	480	595
(再掲)40-74歳※	733	320	144	6.8	90	139	164	217	300	395	498	586	783	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

パントテン酸(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	5.38	1.91	0.370	2.25	2.49	2.87	3.68	5.42	6.61	8.05	8.76	9.76
	30-39歳	58	6.55	2.38	0.351	1.96	2.79	3.54	5.12	6.51	7.84	10.23	11.16	12.61
	40-49歳	96	5.32	1.85	0.184	1.25	2.46	2.78	4.04	5.18	6.48	7.77	8.58	10.81
	50-59歳	84	6.13	1.96	0.216	2.08	3.60	3.87	4.73	5.99	7.00	9.05	9.90	12.55
	60-69歳	115	6.28	1.88	0.182	2.00	3.59	4.08	5.17	6.12	7.58	8.59	9.38	13.35
	70歳以上	168	6.39	2.11	0.244	2.30	3.18	3.70	4.81	6.28	8.10	9.42	9.75	11.30
	(再掲)20歳以上※	563	6.01	2.06	0.118	2.17	2.86	3.45	4.57	5.92	7.30	8.77	9.66	11.44
	(再掲)65-74歳	130	6.62	1.91	0.171	2.52	3.59	4.19	5.46	6.34	7.87	9.22	9.94	13.08
	(再掲)75歳以上	107	6.26	2.07	0.311	2.36	3.20	3.71	4.72	6.25	7.63	9.29	9.60	11.74
	(再掲)20-49歳※	196	5.69	2.10	0.180	1.92	2.64	2.89	4.18	5.67	6.70	8.54	9.54	12.15
(再掲)40-74歳※	356	6.00	1.98	0.118	1.82	3.05	3.63	4.59	5.90	7.29	8.64	9.45	11.20	
女性	20-29歳	40	4.77	1.80	0.334	1.26	1.93	2.45	3.85	4.44	5.76	7.27	8.73	9.55
	30-39歳	87	5.07	1.54	0.215	2.20	2.94	3.15	3.87	4.80	6.10	7.57	7.88	9.29
	40-49歳	85	5.05	1.80	0.257	2.08	2.32	3.03	3.55	4.91	6.19	7.19	8.62	11.12
	50-59歳	97	5.31	1.55	0.189	2.15	2.74	3.16	4.14	5.27	6.34	7.20	7.92	10.96
	60-69歳	145	5.84	2.05	0.167	1.50	3.05	3.55	4.28	5.59	6.94	8.46	9.35	11.99
	70歳以上	187	5.23	1.69	0.114	1.69	2.66	3.33	4.02	5.21	6.09	7.46	7.95	9.73
	(再掲)20歳以上※	641	5.23	1.77	0.087	1.71	2.69	3.15	3.99	5.07	6.28	7.53	8.39	10.79
	(再掲)65-74歳	151	5.70	1.93	0.170	1.58	3.02	3.57	4.24	5.62	6.84	8.30	8.99	11.97
	(再掲)75歳以上	137	5.19	1.76	0.147	1.66	2.47	3.16	3.99	5.13	6.21	7.46	7.86	11.67
	(再掲)20-49歳※	212	4.96	1.72	0.142	1.28	2.43	2.96	3.83	4.83	5.96	7.45	8.05	10.15
(再掲)40-74歳※	377	5.40	1.80	0.115	2.04	2.81	3.25	4.11	5.27	6.38	7.77	8.53	11.36	
男女計	20-29歳	82	5.09	1.87	0.314	1.26	2.40	2.76	3.82	4.90	6.08	7.65	8.65	9.76
	30-39歳	145	5.83	2.14	0.234	2.10	2.91	3.26	4.14	5.61	6.91	8.38	10.21	12.52
	40-49歳	181	5.19	1.82	0.169	1.71	2.47	3.01	3.83	5.03	6.27	7.42	8.58	10.83
	50-59歳	181	5.72	1.81	0.156	2.16	2.90	3.73	4.34	5.55	6.58	8.04	9.21	11.27
	60-69歳	260	6.06	1.98	0.130	1.74	3.21	3.77	4.66	5.91	7.30	8.45	9.36	11.97
	70歳以上	355	5.71	1.96	0.143	1.83	2.90	3.47	4.30	5.44	6.90	8.30	9.31	11.16
	(再掲)20歳以上※	1,204	5.61	1.96	0.091	1.91	2.76	3.25	4.19	5.42	6.70	8.16	9.21	11.16
	(再掲)65-74歳	281	6.13	1.97	0.142	2.15	3.32	3.78	4.68	5.94	7.37	8.53	9.43	11.90
	(再掲)75歳以上	244	5.61	1.95	0.175	1.71	2.74	3.32	4.23	5.39	6.81	8.15	9.25	11.18
	(再掲)20-49歳※	408	5.34	1.96	0.131	1.79	2.47	2.93	3.92	5.12	6.49	7.81	8.86	10.83
(再掲)40-74歳※	733	5.70	1.92	0.100	2.02	2.88	3.44	4.31	5.54	6.70	8.20	9.22	11.22	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

ビタミンC(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	61	48	11	3	11	12	16	54	79	143	178	216
	30-39歳	58	91	54	8	5	13	22	43	81	130	166	201	219
	40-49歳	96	76	44	5	9	16	23	37	75	92	128	172	222
	50-59歳	84	91	59	6	16	24	31	43	79	123	167	202	357
	60-69歳	115	115	73	6	10	19	33	64	103	161	218	272	326
	70歳以上	168	123	73	6	19	36	45	70	112	162	216	288	365
	(再掲)20歳以上※	563	95	65	5	9	16	26	47	80	128	183	216	326
	(再掲)65-74歳	130	128	78	6	12	26	42	67	113	174	232	304	358
	(再掲)75歳以上	107	119	71	8	15	34	46	68	106	161	207	291	363
	(再掲)20-49歳※	196	75	50	6	4	12	16	37	70	96	141	179	218
(再掲)40-74歳※	356	99	65	4	12	22	32	52	84	129	187	221	330	
女性	20-29歳	40	75	52	13	9	14	15	36	69	107	125	189	266
	30-39歳	87	81	55	6	7	19	25	48	73	98	132	197	382
	40-49歳	85	76	44	4	5	23	28	42	74	98	123	158	319
	50-59歳	97	106	60	7	8	30	40	58	101	130	200	232	313
	60-69歳	145	127	84	7	15	24	36	67	111	177	227	314	420
	70歳以上	187	119	73	7	15	32	43	69	109	151	212	240	456
	(再掲)20歳以上※	641	101	69	4	14	19	31	54	86	126	192	223	378
	(再掲)65-74歳	151	132	85	7	14	35	45	71	113	181	229	315	452
	(再掲)75歳以上	137	115	71	8	15	27	41	68	102	148	206	227	414
	(再掲)20-49歳※	212	77	50	5	10	16	22	41	73	100	125	162	280
(再掲)40-74歳※	377	108	71	4	14	25	36	59	93	130	200	237	396	
男女計	20-29歳	82	68	51	10	3	11	15	23	64	92	126	176	266
	30-39歳	145	86	55	5	5	16	25	45	79	116	146	199	297
	40-49歳	181	76	44	4	10	17	27	39	75	94	123	156	220
	50-59歳	181	99	60	6	15	26	36	51	89	130	177	210	307
	60-69歳	260	121	79	5	13	24	35	65	107	161	220	299	398
	70歳以上	355	121	73	6	16	34	45	69	109	157	215	253	370
	(再掲)20歳以上※	1,204	98	67	4	11	16	28	51	84	127	188	220	339
	(再掲)65-74歳	281	130	82	6	14	31	44	71	113	177	230	309	417
	(再掲)75歳以上	244	116	71	7	15	31	42	68	106	152	205	234	369
	(再掲)20-49歳※	408	76	50	5	7	15	17	38	71	98	135	173	221
(再掲)40-74歳※	733	104	69	3	14	25	34	55	89	130	197	231	355	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

食塩相当量(g)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	9.4	4.4	0.73	2.1	2.7	4.2	6.0	9.0	12.5	17.0	18.3	19.6
	30-39歳	58	11.3	4.6	0.57	2.0	4.0	4.9	8.2	10.7	13.5	17.4	20.2	24.3
	40-49歳	96	11.3	3.8	0.59	4.0	5.3	6.7	8.3	11.5	13.2	16.1	17.9	26.2
	50-59歳	84	10.8	3.5	0.54	2.0	5.7	6.9	8.6	10.2	12.3	16.1	17.7	20.6
	60-69歳	115	11.9	4.3	0.44	4.4	5.8	6.7	9.0	11.7	14.2	16.4	19.5	29.3
	70歳以上	168	11.9	4.1	0.38	3.8	6.4	7.0	9.5	11.5	13.7	17.6	19.3	24.4
	(再掲)20歳以上※	563	11.2	4.2	0.23	2.5	5.2	6.1	8.3	10.8	13.2	16.8	18.9	24.2
	(再掲)65-74歳	130	12.3	4.4	0.53	5.6	6.4	7.2	9.1	11.9	14.4	17.3	20.7	28.5
	(再掲)75歳以上	107	11.6	4.0	0.40	3.6	6.5	6.9	9.4	11.0	13.0	17.7	19.5	24.1
	(再掲)20-49歳※	196	10.7	4.3	0.38	2.0	4.2	5.5	7.5	10.6	13.1	16.7	18.0	23.8
(再掲)40-74歳※	356	11.5	4.0	0.31	4.3	5.7	6.9	8.7	11.3	13.5	16.3	18.7	24.4	
女性	20-29歳	40	8.3	3.0	0.52	3.0	3.3	3.8	5.5	8.2	10.5	12.0	13.3	14.0
	30-39歳	87	8.5	2.7	0.32	2.2	4.3	5.2	6.8	7.9	10.3	11.8	14.2	16.5
	40-49歳	85	9.2	3.9	0.40	3.8	4.6	4.6	6.2	8.8	11.0	13.3	17.6	22.3
	50-59歳	97	9.8	3.3	0.33	3.4	4.8	5.2	7.2	10.2	11.9	13.5	15.2	20.4
	60-69歳	145	10.4	4.1	0.43	2.2	5.1	6.0	7.7	10.0	12.2	14.6	18.8	26.4
	70歳以上	187	10.1	3.4	0.24	4.0	5.3	5.9	7.4	10.0	12.1	14.5	16.3	19.9
	(再掲)20歳以上※	641	9.5	3.5	0.19	3.3	4.5	5.3	7.1	9.5	11.6	13.5	15.2	21.4
	(再掲)65-74歳	151	10.5	3.8	0.38	2.3	5.5	6.4	8.0	10.1	12.1	14.6	18.2	24.0
	(再掲)75歳以上	137	10.0	3.5	0.33	3.7	5.1	5.6	7.2	10.0	12.2	14.6	16.2	21.8
	(再掲)20-49歳※	212	8.7	3.3	0.28	3.1	4.3	4.6	6.3	8.2	10.6	12.6	14.1	21.1
(再掲)40-74歳※	377	9.9	3.7	0.23	3.6	4.6	5.5	7.2	9.8	11.9	14.2	15.7	22.0	
男女計	20-29歳	82	8.9	3.8	0.55	2.1	3.3	4.2	5.9	8.5	10.7	13.8	17.1	19.6
	30-39歳	145	9.9	4.1	0.36	2.2	4.4	5.2	7.1	9.7	12.2	15.7	17.4	23.5
	40-49歳	181	10.3	4.0	0.42	4.1	4.6	5.3	7.2	10.1	12.6	15.3	17.8	22.6
	50-59歳	181	10.3	3.4	0.36	3.5	5.0	6.1	7.8	10.2	12.0	14.7	17.4	20.5
	60-69歳	260	11.1	4.3	0.35	3.4	5.7	6.3	8.3	10.7	13.2	15.9	19.4	26.2
	70歳以上	355	10.9	3.8	0.23	3.8	5.5	6.4	8.1	10.8	12.6	15.6	18.0	24.3
	(再掲)20歳以上※	1,204	10.3	4.0	0.18	3.2	4.6	5.6	7.4	10.2	12.4	15.2	17.5	22.9
	(再掲)65-74歳	281	11.4	4.2	0.38	3.7	5.9	6.7	8.5	11.0	13.3	16.2	19.2	25.2
	(再掲)75歳以上	244	10.6	3.8	0.25	3.7	5.3	6.1	7.7	10.6	12.6	15.0	18.0	23.4
	(再掲)20-49歳※	408	9.7	4.0	0.30	2.4	4.3	4.6	6.9	9.5	12.0	14.7	17.3	21.9
(再掲)40-74歳※	733	10.7	3.9	0.24	3.9	5.2	6.1	7.9	10.4	12.6	15.6	17.8	24.2	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

カリウム(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	1,980	917	116	785	822	965	1,381	1,737	2,318	3,267	4,346	4,865
	30-39歳	58	2,581	955	118	682	1,066	1,321	1,947	2,533	3,010	4,094	4,661	4,777
	40-49歳	96	2,185	745	75	542	1,092	1,223	1,678	2,118	2,658	3,350	3,524	4,360
	50-59歳	84	2,409	910	115	720	1,273	1,545	1,745	2,163	2,869	3,834	4,482	5,050
	60-69歳	115	2,715	983	88	556	1,428	1,654	1,989	2,528	3,320	3,958	4,466	6,657
	70歳以上	168	2,820	1,029	108	908	1,408	1,618	2,050	2,661	3,493	4,285	4,809	5,370
	(再掲)20歳以上※	563	2,470	976	58	785	1,118	1,371	1,796	2,306	2,988	3,861	4,361	5,195
	(再掲)65-74歳	130	2,921	1,005	84	1,255	1,512	1,736	2,180	2,850	3,517	4,304	4,684	6,465
	(再掲)75歳以上	107	2,745	1,038	142	809	1,308	1,577	2,006	2,599	3,151	4,248	5,194	5,354
	(再掲)20-49歳※	196	2,228	896	59	731	972	1,213	1,606	2,101	2,717	3,479	4,146	4,795
(再掲)40-74歳※	356	2,511	940	56	858	1,224	1,502	1,855	2,307	3,095	3,794	4,345	5,091	
女性	20-29歳	40	1,855	733	123	385	587	1,137	1,507	1,713	2,239	2,639	2,848	5,196
	30-39歳	87	2,074	679	94	955	1,147	1,305	1,604	2,017	2,434	3,074	3,381	4,324
	40-49歳	85	2,098	765	107	853	1,177	1,315	1,525	1,933	2,584	3,230	3,706	4,289
	50-59歳	97	2,339	747	76	893	1,167	1,418	1,688	2,315	2,732	3,188	3,824	4,649
	60-69歳	145	2,787	1,427	142	505	1,166	1,527	1,982	2,507	3,290	3,936	4,303	9,577
	70歳以上	187	2,408	910	73	805	1,110	1,310	1,828	2,298	2,903	3,471	3,900	5,593
	(再掲)20歳以上※	641	2,300	976	49	679	1,134	1,317	1,643	2,200	2,669	3,401	3,800	5,184
	(再掲)65-74歳	151	2,735	1,325	157	540	1,271	1,555	2,037	2,564	3,170	3,807	4,272	9,285
	(再掲)75歳以上	137	2,361	943	92	795	952	1,253	1,802	2,218	2,737	3,471	3,896	6,314
	(再掲)20-49歳※	212	2,010	735	67	428	1,097	1,241	1,535	1,862	2,398	3,029	3,479	4,362
(再掲)40-74歳※	377	2,439	1,055	64	862	1,180	1,383	1,710	2,328	2,978	3,541	3,921	5,928	
男女計	20-29歳	82	1,920	831	86	397	822	1,010	1,472	1,720	2,302	2,729	4,043	5,196
	30-39歳	145	2,332	866	85	712	1,084	1,308	1,729	2,220	2,734	3,581	4,106	4,751
	40-49歳	181	2,142	754	82	821	1,126	1,269	1,601	2,009	2,622	3,262	3,627	4,096
	50-59歳	181	2,374	831	78	893	1,261	1,470	1,728	2,254	2,864	3,651	3,906	4,974
	60-69歳	260	2,752	1,227	93	532	1,320	1,622	1,986	2,519	3,305	3,936	4,426	8,167
	70歳以上	355	2,580	982	78	812	1,201	1,487	1,922	2,459	3,071	3,885	4,487	5,451
	(再掲)20歳以上※	1,204	2,382	979	48	784	1,129	1,343	1,709	2,239	2,867	3,616	4,151	5,196
	(再掲)65-74歳	281	2,822	1,188	108	918	1,387	1,587	2,110	2,630	3,361	4,016	4,588	7,770
	(再掲)75歳以上	244	2,510	997	89	800	1,110	1,391	1,859	2,325	3,002	3,833	4,319	5,556
	(再掲)20-49歳※	408	2,122	828	52	608	1,010	1,230	1,547	1,979	2,549	3,218	3,788	4,742
(再掲)40-74歳※	733	2,475	999	51	874	1,210	1,399	1,805	2,317	2,998	3,652	4,086	5,083	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

カルシウム(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	390	219	38	107	128	188	219	351	466	698	1,028	1,107
	30-39歳	58	511	258	30	81	175	222	299	429	706	858	1,015	1,293
	40-49歳	96	446	215	26	72	163	195	274	405	587	741	867	1,028
	50-59歳	84	519	293	40	52	202	234	297	455	689	937	1,092	1,600
	60-69歳	115	558	251	29	135	211	258	369	518	719	907	983	1,524
	70歳以上	168	613	285	26	185	226	270	386	562	796	1,050	1,125	1,370
	(再掲)20歳以上※	563	513	266	16	109	188	218	310	463	663	898	1,032	1,338
	(再掲)65-74歳	130	595	258	26	177	229	268	409	567	741	930	1,107	1,461
	(再掲)75歳以上	107	617	298	37	137	210	261	385	548	802	1,073	1,125	1,380
	(再掲)20-49歳※	196	446	233	22	95	173	195	272	400	578	775	974	1,097
(再掲)40-74歳※	356	521	259	16	128	190	233	321	481	663	888	995	1,339	
女性	20-29歳	40	391	203	38	73	129	176	218	338	522	702	824	898
	30-39歳	87	443	198	32	112	150	211	311	425	581	692	853	968
	40-49歳	85	491	304	33	152	178	198	280	426	608	866	1,047	1,744
	50-59歳	97	531	251	28	113	216	265	350	485	687	821	935	1,569
	60-69歳	145	616	472	53	78	194	262	377	547	742	955	1,093	3,025
	70歳以上	187	535	232	19	140	190	247	343	527	664	880	946	1,176
	(再掲)20歳以上※	641	511	300	14	105	178	214	312	467	645	843	953	1,462
	(再掲)65-74歳	151	602	440	53	84	194	263	373	549	728	924	983	2,852
	(再掲)75歳以上	137	530	235	20	128	186	232	342	522	672	853	971	1,176
	(再掲)20-49歳※	212	444	248	23	78	170	187	277	401	559	757	895	1,501
(再掲)40-74歳※	377	548	348	20	145	195	235	332	481	672	901	985	1,679	
男女計	20-29歳	82	390	210	30	74	145	178	226	339	471	694	820	1,107
	30-39歳	145	478	232	25	89	175	216	306	428	603	831	918	1,155
	40-49歳	181	468	263	25	146	171	197	277	420	591	780	969	1,573
	50-59歳	181	525	272	31	88	214	248	312	481	688	891	1,030	1,562
	60-69歳	260	587	380	25	118	210	260	370	529	725	928	1,003	2,128
	70歳以上	355	567	258	19	156	200	259	370	549	705	935	1,087	1,318
	(再掲)20歳以上※	1,204	512	284	13	110	182	215	311	464	655	869	992	1,348
	(再掲)65-74歳	281	599	366	29	145	216	268	379	554	734	928	997	1,905
	(再掲)75歳以上	244	564	264	22	134	195	250	361	533	706	942	1,089	1,354
	(再掲)20-49歳※	408	445	240	20	90	171	194	273	401	572	766	895	1,248
(再掲)40-74歳※	733	535	307	14	142	194	235	327	481	669	893	992	1,510	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

マグネシウム(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	209	78	12.4	83	110	117	147	188	279	307	379	418
	30-39歳	58	278	101	11.1	82	141	159	208	267	330	441	493	522
	40-49歳	96	242	76	10.4	64	136	151	195	231	271	355	380	528
	50-59歳	84	274	103	12.1	78	139	159	200	260	312	437	471	589
	60-69歳	115	290	100	8.4	72	168	190	228	281	331	391	417	853
	70歳以上	168	286	91	10.0	111	145	171	224	277	338	405	458	542
	(再掲)20歳以上※	563	264	96	5.5	84	129	154	200	256	310	392	438	532
	(再掲)65-74歳	130	300	102	8.9	126	173	199	234	293	344	408	429	799
	(再掲)75歳以上	107	279	89	14.2	110	145	167	224	273	328	400	453	521
	(再掲)20-49歳※	196	241	89	7.1	82	114	133	182	226	292	373	421	519
(再掲)40-74歳※	356	273	95	5.8	84	143	173	214	259	320	393	437	552	
女性	20-29歳	40	195	66	11.5	37	69	124	149	192	244	263	298	401
	30-39歳	87	210	67	9.5	88	106	129	157	205	260	297	320	429
	40-49歳	85	228	81	10.6	104	126	147	170	204	263	343	422	497
	50-59歳	97	241	73	7.0	96	127	143	185	243	288	332	367	481
	60-69歳	145	283	161	16.8	59	131	154	196	264	325	373	447	1,120
	70歳以上	187	242	79	6.3	92	116	143	193	238	281	337	382	501
	(再掲)20歳以上※	641	236	98	4.7	86	119	143	177	229	280	334	376	501
	(再掲)65-74歳	151	274	147	17.6	72	129	155	195	258	305	367	424	1,038
	(再掲)75歳以上	137	239	80	7.1	90	112	142	190	231	279	337	380	516
	(再掲)20-49歳※	212	212	73	7.3	50	107	136	164	201	255	303	344	427
(再掲)40-74歳※	377	252	112	6.3	101	131	151	185	240	296	343	413	580	
男女計	20-29歳	82	202	73	8.7	40	95	120	149	190	255	296	328	418
	30-39歳	145	245	92	8.4	83	113	141	182	232	289	373	440	510
	40-49歳	181	235	78	9.6	93	133	148	181	225	271	343	381	505
	50-59歳	181	258	91	7.8	93	136	155	196	247	297	391	446	552
	60-69歳	260	286	134	10.4	72	150	173	218	271	329	388	422	1,060
	70歳以上	355	260	87	6.9	96	126	146	204	252	304	367	418	521
	(再掲)20歳以上※	1,204	250	98	4.6	85	126	145	187	239	294	361	416	522
	(再掲)65-74歳	281	286	129	11.0	106	144	173	216	273	331	401	424	1,033
	(再掲)75歳以上	244	254	86	8.2	93	119	145	199	247	296	362	417	518
	(再掲)20-49歳※	408	227	83	6.2	71	114	136	170	214	270	330	390	497
(再掲)40-74歳※	733	262	105	4.9	100	137	156	199	248	305	375	422	549	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

リン(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	937	285	41	415	459	499	771	915	1,175	1,298	1,435	1,457
	30-39歳	58	1,149	394	44	401	476	649	876	1,127	1,395	1,739	1,927	2,111
	40-49歳	96	983	294	35	326	567	592	787	975	1,147	1,347	1,434	1,988
	50-59歳	84	1,094	352	44	300	654	679	860	1,064	1,265	1,583	1,753	2,242
	60-69歳	115	1,149	332	27	376	667	763	905	1,128	1,353	1,528	1,641	2,540
	70歳以上	168	1,126	335	33	426	557	683	878	1,140	1,354	1,597	1,682	1,832
	(再掲)20歳以上※	563	1,074	340	17	414	553	654	826	1,075	1,283	1,516	1,666	1,950
	(再掲)65-74歳	130	1,189	340	30	505	700	772	929	1,168	1,382	1,638	1,716	2,424
	(再掲)75歳以上	107	1,110	332	43	417	550	637	878	1,101	1,353	1,537	1,647	2,093
	(再掲)20-49歳※	196	1,015	333	24	407	486	586	794	987	1,209	1,410	1,662	1,983
(再掲)40-74歳※	356	1,086	334	19	354	596	680	840	1,079	1,293	1,521	1,674	1,988	
女性	20-29歳	40	863	264	44	229	428	458	685	841	998	1,221	1,291	1,561
	30-39歳	87	867	257	37	360	497	539	680	861	1,065	1,206	1,361	1,501
	40-49歳	85	914	323	37	403	490	571	679	876	1,073	1,268	1,630	2,088
	50-59歳	97	969	280	35	412	505	619	768	966	1,123	1,321	1,526	1,850
	60-69歳	145	1,031	355	27	243	533	631	766	1,017	1,232	1,479	1,622	2,175
	70歳以上	187	958	298	26	347	480	595	736	945	1,129	1,327	1,446	1,801
	(再掲)20歳以上※	641	941	306	13	358	490	582	722	920	1,114	1,300	1,471	1,932
	(再掲)65-74歳	151	1,033	339	33	281	505	655	773	1,014	1,231	1,477	1,624	2,099
	(再掲)75歳以上	137	942	300	26	347	475	549	725	935	1,115	1,335	1,440	1,825
	(再掲)20-49歳※	212	883	286	20	354	458	548	683	855	1,059	1,230	1,338	1,953
(再掲)40-74歳※	377	977	322	17	373	510	620	749	947	1,164	1,340	1,605	2,074	
男女計	20-29歳	82	901	276	32	263	443	493	731	868	1,130	1,256	1,315	1,561
	30-39歳	145	1,011	361	34	397	501	552	747	974	1,202	1,506	1,735	2,019
	40-49歳	181	949	310	30	374	531	592	716	918	1,108	1,341	1,464	2,026
	50-59歳	181	1,032	324	33	405	594	631	795	1,004	1,204	1,510	1,609	2,100
	60-69歳	260	1,089	349	22	338	597	681	829	1,083	1,291	1,518	1,622	2,168
	70歳以上	355	1,028	324	25	369	516	613	786	998	1,266	1,463	1,638	1,794
	(再掲)20歳以上※	1,204	1,005	329	13	372	510	604	782	977	1,207	1,417	1,597	1,944
	(再掲)65-74歳	281	1,106	348	27	367	613	699	864	1,084	1,315	1,534	1,685	2,103
	(再掲)75歳以上	244	1,007	323	28	348	500	598	779	982	1,257	1,436	1,580	1,821
	(再掲)20-49歳※	408	951	317	18	392	471	561	726	932	1,153	1,323	1,498	1,959
(再掲)40-74歳※	733	1,031	332	15	371	572	632	787	998	1,231	1,452	1,623	2,035	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

鉄(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	7.0	2.7	0.43	2.4	3.5	3.9	5.1	6.4	8.8	10.2	12.1	16.5
	30-39歳	58	8.6	3.3	0.34	1.4	3.4	4.3	6.3	8.4	10.3	12.7	15.8	16.8
	40-49歳	96	7.4	2.5	0.30	2.0	4.0	4.6	5.7	7.1	8.4	10.5	11.9	15.7
	50-59歳	84	8.4	3.1	0.39	2.1	4.1	5.0	6.4	7.8	10.0	13.1	14.8	19.0
	60-69歳	115	9.0	3.0	0.27	3.3	4.7	5.3	6.6	9.0	10.6	13.0	14.6	19.3
	70歳以上	168	9.1	3.0	0.23	4.1	4.9	5.7	7.0	8.7	11.1	13.2	14.4	17.5
	(再掲)20歳以上※	563	8.3	3.0	0.18	2.5	4.0	4.8	6.2	7.8	10.0	12.6	14.3	16.6
	(再掲)65-74歳	130	9.4	3.2	0.28	3.3	4.7	5.6	7.3	9.3	11.3	13.3	14.6	22.5
	(再掲)75歳以上	107	8.8	2.6	0.30	3.8	4.6	5.4	7.0	8.4	10.4	12.6	13.7	15.0
	(再掲)20-49歳※	196	7.6	2.9	0.22	2.0	3.7	4.2	5.8	7.2	8.9	11.3	12.7	16.6
(再掲)40-74歳※	356	8.5	3.1	0.18	3.1	4.4	5.1	6.4	7.8	10.1	12.9	14.4	17.8	
女性	20-29歳	40	6.4	2.5	0.46	1.6	2.4	4.1	5.1	6.2	7.6	9.0	9.3	16.7
	30-39歳	87	6.8	2.2	0.30	2.7	3.4	4.3	5.1	6.4	8.1	9.4	10.4	12.9
	40-49歳	85	6.9	2.7	0.35	2.6	3.6	3.8	5.1	6.5	8.3	11.0	11.9	15.2
	50-59歳	97	7.8	2.5	0.30	2.8	3.9	4.7	6.2	7.5	9.2	11.7	12.6	13.9
	60-69歳	145	9.7	10.1	1.17	1.5	3.7	4.8	6.3	8.4	10.4	12.5	14.4	65.7
	70歳以上	187	7.9	2.8	0.22	2.7	3.9	4.6	6.0	7.6	9.0	11.6	13.3	17.6
	(再掲)20歳以上※	641	7.7	4.9	0.20	2.5	3.6	4.4	5.6	7.2	8.9	11.1	12.8	17.0
	(再掲)65-74歳	151	9.3	9.3	1.05	2.7	4.1	5.1	6.5	7.8	9.8	11.8	14.0	62.9
	(再掲)75歳以上	137	7.9	3.0	0.28	2.5	3.7	4.5	6.0	7.5	9.3	12.3	13.8	18.4
	(再掲)20-49歳※	212	6.7	2.5	0.23	1.8	3.5	3.8	5.1	6.4	8.0	9.3	11.0	15.3
(再掲)40-74歳※	377	8.1	6.1	0.36	2.7	3.8	4.4	5.7	7.4	9.3	11.6	13.0	18.8	
男女計	20-29歳	82	6.7	2.6	0.34	1.7	3.1	3.9	5.1	6.2	8.0	10.0	11.3	16.7
	30-39歳	145	7.7	2.9	0.26	1.7	3.5	4.3	5.7	7.5	9.1	11.3	12.9	16.7
	40-49歳	181	7.1	2.6	0.30	2.5	3.7	4.1	5.5	6.8	8.4	10.6	11.9	15.6
	50-59歳	181	8.1	2.8	0.30	2.5	4.0	4.9	6.4	7.7	9.5	12.5	13.3	17.7
	60-69歳	260	9.4	7.5	0.62	3.1	4.1	5.1	6.5	8.7	10.4	12.9	14.4	39.4
	70歳以上	355	8.4	2.9	0.19	3.2	4.4	5.0	6.5	7.9	9.9	12.5	14.2	17.6
	(再掲)20歳以上※	1,204	8.0	4.1	0.15	2.5	3.8	4.5	5.9	7.5	9.4	11.9	13.4	16.7
	(再掲)65-74歳	281	9.4	7.1	0.59	3.2	4.5	5.3	6.8	8.4	10.3	13.0	14.4	33.7
	(再掲)75歳以上	244	8.3	2.9	0.23	2.8	4.0	4.7	6.4	7.9	9.8	12.5	13.7	16.6
	(再掲)20-49歳※	408	7.1	2.7	0.18	1.9	3.6	4.1	5.3	6.7	8.6	10.4	12.1	16.5
(再掲)40-74歳※	733	8.3	4.8	0.20	2.9	3.9	4.8	6.0	7.7	9.7	12.1	13.9	17.5	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

亜鉛(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	8.0	2.6	0.33	3.5	3.6	4.6	6.0	7.7	10.1	12.1	12.8	13.2
	30-39歳	58	9.9	3.5	0.46	2.5	4.3	5.2	7.2	9.7	12.5	15.3	16.1	19.4
	40-49歳	96	8.5	2.5	0.24	2.8	4.9	5.5	6.6	8.2	9.9	12.1	13.9	15.4
	50-59歳	84	9.0	2.5	0.24	3.7	6.2	6.4	7.2	8.6	10.3	13.1	13.5	17.1
	60-69歳	115	9.2	2.6	0.22	4.0	5.5	6.2	7.5	8.9	10.5	12.2	14.0	20.7
	70歳以上	168	8.8	2.4	0.25	3.3	4.8	5.8	6.9	8.8	10.5	12.0	12.8	14.4
	(再掲)20歳以上※	563	8.9	2.7	0.12	3.5	4.9	5.5	6.9	8.7	10.4	12.6	13.3	16.2
	(再掲)65-74歳	130	9.4	2.6	0.21	4.0	5.4	6.4	7.7	9.1	10.6	12.3	13.9	20.1
	(再掲)75歳以上	107	8.6	2.4	0.33	3.2	4.7	5.7	6.7	8.6	10.3	11.7	12.8	14.8
	(再掲)20-49歳※	196	8.7	3.0	0.19	3.0	4.5	5.1	6.5	8.5	10.4	12.7	14.0	17.6
(再掲)40-74歳※	356	8.9	2.5	0.12	3.6	5.2	6.1	7.1	8.7	10.3	12.2	13.5	15.8	
女性	20-29歳	40	6.5	2.1	0.30	1.6	3.0	4.1	4.6	6.3	8.0	9.5	10.4	11.9
	30-39歳	87	7.0	2.1	0.28	2.2	3.6	4.1	5.8	6.9	8.3	9.6	9.9	14.4
	40-49歳	85	7.4	2.1	0.28	3.4	4.1	4.9	6.0	7.0	8.8	10.0	11.9	14.1
	50-59歳	97	7.3	2.1	0.25	2.9	4.1	4.8	6.0	7.3	8.3	9.6	10.6	16.9
	60-69歳	145	7.6	2.6	0.19	2.1	3.9	4.5	6.0	7.2	9.0	11.0	13.1	14.7
	70歳以上	187	7.1	2.0	0.14	2.8	3.9	4.4	5.8	6.9	8.5	9.6	10.1	12.8
	(再掲)20歳以上※	641	7.2	2.2	0.10	2.9	3.9	4.5	5.8	6.9	8.5	9.8	10.9	13.6
	(再掲)65-74歳	151	7.6	2.3	0.17	2.4	3.9	4.6	6.0	7.5	9.0	10.1	11.7	14.6
	(再掲)75歳以上	137	6.9	2.0	0.18	2.7	3.7	4.4	5.7	6.8	8.1	9.7	10.3	13.2
	(再掲)20-49歳※	212	7.0	2.1	0.17	2.4	3.7	4.4	5.4	6.8	8.3	9.5	10.4	12.8
(再掲)40-74歳※	377	7.4	2.2	0.12	2.9	4.1	4.8	6.0	7.2	8.8	10.0	11.5	14.3	
男女計	20-29歳	82	7.2	2.5	0.25	1.9	3.6	4.5	5.3	7.0	8.9	10.9	12.2	13.2
	30-39歳	145	8.5	3.2	0.32	2.4	3.9	4.9	6.3	8.0	10.0	12.8	15.2	18.8
	40-49歳	181	7.9	2.4	0.21	3.4	4.5	5.1	6.2	7.5	9.5	11.1	12.6	14.7
	50-59歳	181	8.2	2.4	0.20	3.1	4.6	5.6	6.5	7.9	9.4	11.4	13.2	17.1
	60-69歳	260	8.4	2.7	0.17	2.9	4.4	5.4	6.7	8.3	10.0	11.6	13.1	16.3
	70歳以上	355	7.8	2.4	0.16	2.9	4.0	4.8	6.1	7.5	9.4	10.9	12.1	13.8
	(再掲)20歳以上※	1,204	8.0	2.6	0.10	3.0	4.1	4.9	6.2	7.7	9.5	11.4	12.8	15.4
	(再掲)65-74歳	281	8.4	2.6	0.15	2.9	4.6	5.5	6.7	8.4	9.7	11.6	12.9	15.9
	(再掲)75歳以上	244	7.6	2.4	0.20	3.0	3.9	4.5	6.0	7.2	9.2	10.7	11.9	13.9
	(再掲)20-49歳※	408	7.9	2.7	0.16	2.8	4.0	4.6	6.0	7.5	9.5	11.9	12.8	15.9
(再掲)40-74歳※	733	8.2	2.5	0.10	3.3	4.5	5.3	6.5	7.9	9.6	11.5	12.8	15.1	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。

エネルギー及び栄養素の摂取量(20歳以上性年齢階級別)

銅(mg)

性別	年齢区分	人数	平均値	標準偏差	標準誤差	パーセンタイル								
						1	5	10	25	50	75	90	95	99
男性	20-29歳	42	1.08	0.33	0.051	0.43	0.59	0.68	0.85	1.01	1.34	1.60	1.74	1.78
	30-39歳	58	1.39	0.50	0.063	0.35	0.51	0.72	1.14	1.37	1.66	1.98	2.24	3.11
	40-49歳	96	1.15	0.32	0.039	0.42	0.67	0.73	0.88	1.11	1.36	1.62	1.72	1.99
	50-59歳	84	1.31	0.40	0.055	0.47	0.77	0.83	1.05	1.23	1.50	1.99	2.06	2.55
	60-69歳	115	1.44	0.52	0.060	0.44	0.79	0.94	1.15	1.35	1.64	1.92	2.31	3.50
	70歳以上	168	1.37	0.41	0.039	0.56	0.74	0.90	1.09	1.34	1.59	1.92	2.10	2.66
	(再掲)20歳以上※	563	1.29	0.44	0.021	0.42	0.68	0.80	1.00	1.26	1.50	1.84	1.98	2.70
	(再掲)65-74歳	130	1.48	0.52	0.052	0.60	0.89	0.99	1.17	1.40	1.66	1.97	2.51	3.50
	(再掲)75歳以上	107	1.33	0.39	0.045	0.56	0.76	0.88	1.07	1.30	1.50	1.90	2.07	2.63
	(再掲)20-49歳※	196	1.19	0.40	0.026	0.39	0.60	0.70	0.89	1.18	1.42	1.73	1.95	2.32
(再掲)40-74歳※	356	1.32	0.44	0.024	0.47	0.73	0.83	1.04	1.28	1.53	1.85	2.00	3.40	
女性	20-29歳	40	0.94	0.32	0.061	0.16	0.38	0.56	0.69	0.92	1.17	1.31	1.53	1.91
	30-39歳	87	1.00	0.30	0.037	0.42	0.53	0.62	0.79	0.97	1.18	1.40	1.61	2.08
	40-49歳	85	1.06	0.31	0.046	0.51	0.57	0.68	0.84	1.02	1.26	1.48	1.70	1.96
	50-59歳	97	1.09	0.33	0.039	0.33	0.62	0.67	0.87	1.06	1.27	1.45	1.64	2.59
	60-69歳	145	1.19	0.38	0.033	0.35	0.61	0.76	0.91	1.15	1.39	1.71	1.89	2.34
	70歳以上	187	1.11	0.32	0.024	0.46	0.66	0.69	0.89	1.10	1.30	1.51	1.71	2.13
	(再掲)20歳以上※	641	1.08	0.34	0.019	0.35	0.57	0.68	0.85	1.06	1.27	1.50	1.68	2.02
	(再掲)65-74歳	151	1.17	0.35	0.030	0.35	0.66	0.77	0.92	1.16	1.34	1.61	1.88	2.15
	(再掲)75歳以上	137	1.10	0.33	0.031	0.46	0.64	0.68	0.89	1.06	1.30	1.52	1.69	2.27
	(再掲)20-49歳※	212	1.00	0.32	0.034	0.27	0.55	0.64	0.78	0.96	1.20	1.40	1.59	1.89
(再掲)40-74歳※	377	1.12	0.34	0.019	0.45	0.62	0.70	0.88	1.09	1.31	1.54	1.78	2.11	
男女計	20-29歳	82	1.01	0.33	0.045	0.19	0.53	0.63	0.78	0.99	1.24	1.46	1.69	1.91
	30-39歳	145	1.20	0.45	0.042	0.37	0.53	0.68	0.89	1.17	1.45	1.84	1.98	2.74
	40-49歳	181	1.10	0.32	0.035	0.50	0.64	0.70	0.87	1.06	1.33	1.56	1.69	1.94
	50-59歳	181	1.20	0.38	0.041	0.47	0.67	0.77	0.97	1.13	1.38	1.73	1.99	2.54
	60-69歳	260	1.31	0.47	0.036	0.37	0.72	0.83	1.04	1.26	1.52	1.83	1.94	3.50
	70歳以上	355	1.22	0.38	0.028	0.46	0.67	0.76	0.96	1.18	1.41	1.74	1.92	2.40
	(再掲)20歳以上※	1,204	1.18	0.40	0.017	0.43	0.63	0.71	0.90	1.14	1.39	1.68	1.91	2.36
	(再掲)65-74歳	281	1.32	0.47	0.033	0.39	0.69	0.81	1.05	1.26	1.52	1.84	2.00	3.50
	(再掲)75歳以上	244	1.19	0.37	0.031	0.46	0.67	0.71	0.93	1.16	1.38	1.65	1.90	2.34
	(再掲)20-49歳※	408	1.10	0.37	0.024	0.37	0.56	0.68	0.84	1.06	1.33	1.59	1.80	2.15
(再掲)40-74歳※	733	1.22	0.41	0.019	0.47	0.67	0.77	0.94	1.17	1.41	1.71	1.92	2.53	

注) 値は保健所圏域ごとの標本数の偏りを調整する重み付けをして算出。

※保健所圏域ごとの標本数の偏りを調整する重み付けと県の実際のH28人口構成に合わせるための重み付けをした値。